AGRI MAGAZINE
(International E-Magazine for Agricultural Articles)
Volume: 02, Issue: 11 (November, 2025)
Available online at http://www.agrimagazine.in
©Agri Magazine, ISSN: 3048-8656

Maize Seed Processing: Steps, Standards, and Quality Control

*Merajul Islam
M.Sc. (Genetics and Plant Breeding), University of Calcutta, Kolkata, India
*Corresponding Author's email: merajul54997@gmail.com

Seed processing is one of the most critical steps in ensuring high-quality seed production for any crop, particularly for hybrid and certified maize (*Zea mays* L.) seed. The primary objective of seed processing is to improve the physical purity, germination capacity, vigor, and storability of the seed before it reaches the farmer. Maize, being a cross-pollinated crop, often shows genetic and physical variation in harvested lots. Thus, scientific and standardized processing ensures that only healthy, pure, and uniform seeds are marketed for sowing. A well-designed processing system maintains seed viability (≥85%), moisture content (≤13%), and ensures freedom from inert matter, weed seeds, and diseased kernels.

Steps in Maize Seed Processing

a) Receiving and Initial Inspection

The first step involves receiving the harvested seed lots from production fields along with field tags, grower details, and field history. Each lot is inspected visually for mixtures, off-types, diseased cobs, and improper drying.

Samples are collected and tested for moisture content, purity, and germination to decide further processing. Lots with severe admixture or low quality are rejected at this stage.

b) Pre-Drying

Freshly harvested maize seed usually has a high moisture content (18–25%), which is unsuitable for shelling and storage. Therefore, pre-drying is done to reduce the moisture to about 12–13%.

Drying can be done by:

Sun drying under shade with frequent turning, or Mechanical drying using forced-air dryers at temperatures below 40°C to prevent embryo damage. Proper drying ensures safe shelling and minimizes mechanical injury.

c) Shelling

After drying, cobs are shelled using maize shellers or mechanical shelling machines.

During shelling:

- Avoid damage to the embryo and seed coat.
- Maintain individual lot identity.
- Remove broken, chaffy, and underdeveloped kernels.

d) Cleaning and Grading

This is a crucial step for removing inert matter, dust, diseased seeds, and off-sized kernels. A sequence of machines is used:

Primary Cleaning

- Purpose: To remove large, small, and light impurities from the seed lot after pre-cleaning.
- Outcome: Seeds become relatively pure and uniform in size — suitable for further precision cleaning.

Secondary Cleaning (Fine Cleaning)

- Purpose: To achieve higher seed purity and uniformity by removing closely similar inert or damaged materials that escaped the primary cleaner.

- Outcome: Seed lot achieves purity above 98–99%, meeting certification standards.

Gravity Separation

- Purpose: To separate seeds based on specific gravity (density) — removing immature, shriveled, insect-damaged, or low-density seeds.

Machine Used:

- Gravity Separator:

✓ Works on the principle of air flow and vibration.

✓ Heavier (high-quality) seeds move upward; lighter (inferior) seeds move downward and are discarded.

- Outcome: Only physiologically mature, dense, and viable maize seeds are retained for final treatment and packaging.

The cleaned and graded seed should meet the physical purity requirement of ≥98%.

e) Seed Treatment

Before packaging, processed seeds are treated with fungicides and polymers to prevent disease infection and enhance seed appearance and flowability.

Maize Seed Treatment: **Chemicals and Their Description (Per 20 Tons of Seed)**

Chemical	Dose	Function/Description
Polymer	40 kg	Used as a protective coating material to form a thin film around the seed surface. Improves seed appearance, flowability, and reduces dust-off during handling. Also ensures uniform adhesion of fungicides and colourants.
Yashika (binder/coating agent)	10 kg	Acts as a binder that helps chemicals (fungicides, colour, and polymers) adhere firmly to the seed surface. Prevents chemical loss and uneven coating during mechanical mixing.
Food Colour	2 Kg	Added for seed identification, branding, and safety marking (treated seeds should be easily distinguishable). Commonly edible or non-toxic synthetic colours are used.
Seed Star (Sticker/Adhesive)	10 Kg	A sticking or adhesive agent that enhances the attachment of seed treatment chemicals (especially fungicides and polymers) to the seed coat, improving treatment uniformity and durability.
Metalaxyl (Fungicide) (Brand names: Apron, Ridomil)	15 Kg	A systemic fungicide effective against downy mildew, Pythium, and Phytophthora diseases. Protects seeds and seedlings during early germination. Mode of Action: Inhibits RNA synthesis in fungi. Chemical group: Phenylamide.
Carrier Water	200 liters	

Fig. 1- Corn seed processing plant

Fig.2- Treated Seeds

Purpose of Seed Treatment in Maize

- Protects seeds from seed-borne and soil-borne pathogens.
- Improves seed longevity and vigor.
- Enhances early germination and field emergence.
- Ensures safe and identifiable treated seeds before marketing.

f) Packaging and Labeling

Proper packaging maintains seed quality during transport and storage. Common materials include: HDPE bags, laminated pouches, or cloth bags (5–20 kg capacity).

Each bag is labeled with:

- Crop, variety, lot number
- Germination %, purity %, and moisture %
- Month/year of testing and packaging
- Certification tag (if applicable)

Correct packaging prevents moisture absorption and mechanical injury.

Quality Control Parameters

At every stage, seed quality control ensures that final lots meet national and international seed standards. Key parameters are:

Parameter	Standard Value (Certified Seed)	Testing Method
Physical Purity	≥ 98%	Physical purity test
Germination of hybrid seed	≥ 90%	Germination paper or sand test
Moisture Content	≤ 13%	Digital moisture meter / oven method
Inert Matter	≤ 2%	Physical separation
Seed Health	Free from major fungal or bacterial pathogens	Seed health test

Quality assurance is carried out by Seed Testing Laboratories (STLs) under State Seed Certification Agencies (SSCA).

Storage and Distribution

Processed maize seed should be stored under cool, dry, and aerated conditions.

- Optimum storage temperature: 10–15°C
- Relative humidity: ≤40%
- Stack bags on wooden pallets to prevent moisture absorption.

Regular monitoring of moisture and germination is essential. Seeds must be distributed before viability drops below the minimum standard.

Conclusion

Scientific maize seed processing plays a crucial role in ensuring seed purity, vigor, and long-term storability. By adopting modern equipment like gravity separators, color sorters, and automatic seed treaters, seed industries can maintain international standards. Proper processing not only enhances the market value of the seed but also contributes to higher crop productivity and farmer satisfaction. At the least I can say, from field to bag, every stage of maize seed processing adds precision, protection, and performance to the final product.

References

1. International Seed Testing Association (ISTA), 2022. ISTA Rules for Seed Testing.
2. FAO, 2020. Seed Processing and Quality Control Guidelines.
3. Indian Minimum Seed Certification Standards (IMSCS), 2013.
4. Basu, A.K. (2019). Seed Science and Technology. PHI Learning Pvt. Ltd., New Delhi.