AGRI MAGAZINE
(International E-Magazine for Agricultural Articles)
Volume: 02, Issue: 12 (December, 2025)
Available online at http://www.agrimagazine.in

Major Insect Pests of Carrot and Their Management

*Purushotam Sharma[1], Madan Lal Choudhary[2], R P Ghasolia[3], D L Bagri[4], Ratan Lal Sharma[5] and D. K. Bairwa[6]
[1]Asst. Professor (Entomology), Sri Karan Narendra College of Agriculture, Jobner, India
[2]Assoc. Professor (Horticulture), Sri Karan Narendra College of Agriculture, Jobner, India
[3]Professor (Plant Pathology), Sri Karan Narendra College of Agriculture, Jobner, India
[4]Asst. Prof. (Plant Physiology), Sri Karan Narendra College of Agriculture, Jobner, India
[5]Assistant Professor (Plant Pathology), Agriculture Research Station, Jalore, India
[6]Assistant Professor (Entomology), College of Agriculture, Kotputli, India
*Corresponding Author's email: psharma.ento@sknau.ac.in

Carrots (*Daucus carota* L.) are attacked by several insect pests throughout the growing season. The most economically important pests worldwide are the **carrot fly (*Psila rosae*)**, **carrot weevil (*Listronotus oregonensis*)**, and various species of **aphids** (especially the willow-carrot aphid, *Cavariella aegopodii*). Other occasional but locally serious pests include cutworms, wireworms, leafhoppers, and armyworms. Below is a detailed overview of the major pests, their damage symptoms, life cycle highlights, and integrated management strategies.

Pest	Scientific Name	Primary Damage	Key Identification / Life Cycle Notes	Management Strategies
Carrot Fly	*Psila rosae*	Rusty brown tunnels in roots (maggots)	Small black fly with yellowish head; 2–3 generations/year; adults emerge Apr–May and again Aug–Sep (temperate regions)	• Row covers / insect netting (from sowing) • Delay sowing until early June (to miss first generation) • Crop rotation (avoid Umbelliferae for 3–4 years) • Vertical barriers/fences (60–90 cm high) • Entomopathogenic nematodes (Steinernema feltiae) against larvae • Spinosad or cyantraniliprole (where registered)
Carrot Weevil	*Listronotus oregonensis*	Serpentine mines in crown & upper root	Small (5 mm) greyish-black snout beetle; overwinters as adult; 1 generation/year in northern areas	• Deep plowing in fall to bury overwintering adults • Crop rotation (3+ years) • Destroy volunteer carrots & wild hosts • Border sprays with pyrethroids or neonicotinoids early season • Trap crops (early-planted carrots around main field)

Willow-Carrot Aphid	*Cavariella aegopodii*	Distorted foliage, stunting; virus vector (Carrot motley dwarf virus complex)	Small greenish aphid with white waxy fluff; migrates from willow trees in spring	• Reflective mulches (silver plastic) to repel alates • Mineral oil + insecticidal soap sprays • Systemic neonicotinoids (imidacloprid, clothianidin) as seed treatment or drench (note pollinator concerns) • Natural enemies: lady beetles, syrphid flies, parasitoids
Cutworms (several spp.)	Noctuidae (*Agrotis*, *Peridroma*, etc.)	Seedlings cut at soil level at night	Large grey/brown caterpillars that curl into "C" when disturbed	• Weed-free fields before planting • Bait formulations with *Bacillus thuringiensis var. kurstaki* (Bt) or spinosad • Protective collars around transplants
Wireworms	*Agriotes spp.*, *Limonius spp.*	Small round holes and tunnels in roots	Thin, hard, yellow-brown larvae; multi-year life cycle in soil	• Long crop rotation (avoid grasses and cereals) • Soil incorporation of entomopathogenic nematodes or fungi (*Metarhizium*) • Seed treatments with neonicotinoids or diamides where permitted
Aster Leafhopper		Aster yellows phytoplasma transmission	Small wedge-shaped green insect	• Insect netting • Insecticide sprays timed to migration periods • Remove infected plants quickly

Integrated Pest Management (IPM) Recommendations for Carrot

1. **Cultural Control (most important foundation)**

- Practice 3–4 year crop rotation away from Apiaceae.
- Use raised beds and good drainage to reduce carrot fly oviposition.
- Sow later (after mid-May in many regions) to avoid peak carrot fly flights.
- Remove crop residues and volunteer carrots promptly.

2. **Physical Barriers**

- Fine-mesh insect netting or fleece (17–21 g/m²) from emergence until harvest is the single most effective non-chemical method against carrot fly and aphids.

3. **Monitoring & Thresholds**

- Yellow sticky traps or Rebell® orange traps for carrot fly adults.
- Pheromone traps for carrot weevil.
- Scout weekly for aphids and leafhopper adults.

4. **Biological Control**

- *Steinernema feltiae* and *S. carpocapsae* nematodes for carrot fly larvae and carrot weevil larvae.
- Release of predatory bugs (*Orius*, lady beetles) and parasitoids where commercially available.

5. **Chemical Control (use judiciously)**

- Seed treatments (cyantraniliprole, clothianidin) give season-long protection against early pests.
- Foliar options: spinosad, spinetoram, lambda-cyhalothrin, acetamiprid, flupyradifurone.
- Always follow local registration, pre-harvest intervals (PHI), and resistance-management guidelines.

6. **Resistant/Tolerant Varieties**

- Partial resistance to carrot fly: 'Flyaway', 'Resista fly', 'Sytan', 'Maestro'.
- No strong resistance to carrot weevil yet, but early-maturing varieties can escape severe damage.

Conclusion

The carrot fly remains the number one pest in most temperate carrot-growing regions, but a combination of insect netting, delayed sowing, and crop rotation can reduce damage to negligible levels without insecticides. Carrot weevil is the primary concern in North America and requires vigilant rotation and monitoring. Aphids are best managed preventively with reflective mulches and seed treatments, especially where carrot virus diseases are prevalent. Adopting a well-planned IPM program that emphasizes cultural and physical controls, supported by selective biological and chemical tools when necessary, ensures high-quality carrot production while minimizing environmental impact and pesticide resistance risks.