AGRI MAGAZINE
(International E-Magazine for Agricultural Articles)
Volume: 02, Issue: 12 (December, 2025)
Available online at http://www.agrimagazine.in
©Agri Magazine, ISSN: 3048-8656

Combating Fungal Frenemies: A Guide to Horticulture Pathology

*Dr. Sanjay Kumar Bhariya[1], Pritam[2], Dr. Suchismita Mishra[3], Gajanand Chhimpa[4] and Gursevak Singh[5]
[1]Assistant Professor, Department of Molecular Biology and Biotechnology, College of Agriculture, IGKV, Raipur, Chhattisgarh-492012, India
[2]Department of Plant Pathology, School of Agricultural Sciences, Nagaland University, Nagaland-797106, India
[3]Assistant Professor, Department of Floriculture and Landscaping, Faculty of Agricultural Sciences Siksha 'O' Anusandhan, Bhubaneswer, Odisha-751029, India
[4]Department of Entomology, College of Agriculture, CAU, Imphal, Manipur, India
[5]Department of Entomology, School of Agricultural Sciences, Nagaland University, Nagaland-797106, India
*Corresponding Author's email: sanjay.crin@gmail.com

Horticulture contributes substantially to global food, nutritional, and economic security. However, fungal diseases continue to challenge crop productivity, accounting for nearly 20–30% average yield losses in fruits, vegetables, ornamentals, and nursery crops (Strange and Scott, 2005). Fungi such as *Fusarium*, *Botrytis*, *Sclerotinia*, *Alternaria*, and *Phytophthora* (oomycetes but similar in pathology) exhibit high virulence, rapid spore dissemination, and environmental adaptability. As climate change increases humidity and temperature variability, the incidence and severity of fungal diseases are projected to rise (Chakraborty and Newton, 2011). This article reviews key fungal pathogens and outlines sustainable management strategies.

Major Fungal Pathogens in Horticulture

Fungal plant pathogens are diverse in taxonomy, infection biology, and host range.

1 Soil-borne Pathogens

- *Fusarium oxysporum* – causes vascular wilt through xylem colonization, leading to systemic blockage (Michielse and Rep, 2009).
- *Rhizoctonia solani* – responsible for damping-off and root rots in vegetables and ornamentals.
- *Sclerotinia sclerotiorum* – produces sclerotia enabling long-term survival in soil.

2 Foliar Pathogens

- *Erysiphe* spp. – powdery mildews manifesting as white, powdery mycelium.
- *Alternaria* spp. – major leaf spot pathogens in solanaceous and cruciferous crops.

3 Post-Harvest Pathogens

- *Botrytis cinerea* – gray mold affecting berries, tomatoes, ornamentals.
- *Penicillium* spp. – major pathogens in stored fruits.

4 Oomycetes (Fungus-like Organisms)

Though taxonomically distinct, oomycetes are included due to similar epidemiology:

- *Phytophthora infestans* – causes late blight.
- *Pythium* spp. – induces damping-off in seedlings.

These pathogens cause extensive economic damage and necessitate precise diagnostic and management strategies.

Epidemiology and Disease Development

The classical "disease triangle"—susceptible host, virulent pathogen, and favorable environment—is crucial for pathogenesis (Agrios, 2005).

1 Pathogen Biology

Fungi disseminate through airborne spores, rain splash, soil movement, vectors, or contaminated tools (Garrett *et al.*, 2006). Moisture and relative humidity above 85% typically favor germination and infection.

2 Host Susceptibility

Plant stress due to nutrient deficiency, water imbalance, or pest damage predisposes crops to fungal invasion.

3 Environmental Conditions

High humidity, poor airflow, suboptimal soil drainage, and temperature fluctuations commonly increase disease incidence.

Understanding epidemiological factors aids in developing predictive models and targeted control measures.

Disease Diagnosis and Detection

Accurate and early detection is essential for effective management.

1 Traditional Methods

- Symptom-based identification
- Microscopic observation
- Culture-based pathogen isolation

2 Molecular and Advanced Tools

- **PCR and qPCR** for species-specific detection (Kulik *et al.*, 2020)
- **DNA barcoding** (ITS sequencing)
- **LAMP assays** for field-level rapid detection
- **Imaging and AI tools** for real-time diagnosis (Sunil *et al.*, 2023)

Advancements in molecular diagnostics significantly reduce misidentification and improve management speed.

Integrated Disease Management (IDM)

1 Cultural Practices

- Crop rotation with non-host species
- Use of disease-free planting materials
- Drip irrigation to reduce leaf wetness
- Proper pruning and canopy management
- Soil solarization for pathogen suppression

These reduce inoculum load and break disease cycles.

2 Biological Control

Biocontrol agents provide eco-friendly protection:

- *Trichoderma harzianum* antagonizes soil-borne fungi by mycoparasitism and induced systemic resistance (Harman, 2006).
- *Bacillus subtilis* and *Pseudomonas fluorescens* suppress foliar and root pathogens.
- Mycorrhizal fungi enhance nutrient uptake and plant resilience.

3 Chemical Control

Although fungicides remain widely used, resistance development is a concern.

Major fungicide groups:

- **Protectants:** copper compounds, sulfur
- **Systemics:** triazoles, strobilurins (QoI), carboxamides (SDHI)
- **Contact fungicides:** mancozeb, chlorothalonil

Rotation based on FRAC codes is essential to prevent resistance (FRAC, 2024).

4 Host Resistance and Biotechnology

Breeding for resistance remains the most sustainable strategy. Molecular breeding and gene-editing technologies (e.g., CRISPR-Cas9) show promise for generating durable disease-resistant cultivars (Wang *et al.*, 2016).

Future Directions

Climate change, globalization of trade, and intensive horticulture will likely increase fungal disease pressure. Future priorities include:

- Genome-based pathogen surveillance
- Development of AI-driven early warning systems
- Expanded use of biologicals and RNAi-based fungicides
- Breeding programs focused on multi-pathogen resistance

Greater integration of traditional knowledge and advanced biotechnology is essential for sustainable horticulture.

Conclusion

Fungal pathogens—our "frenemies"—play dual roles in horticulture, contributing to soil health while posing major threats to crop productivity. Effective management requires a holistic understanding of pathogen biology, host–pathogen interactions, environmental influences, and emerging diagnostic and control strategies. Integrated Disease Management offers a sustainable framework to minimize losses and build resilient horticultural systems.

References

1. Agrios, G. N. (2005). *Plant Pathology* (5th ed.). *Academic Press.*
2. Chakraborty, S., and Newton, A. C. (2011). Climate change, plant diseases and food security: An overview. *Plant Pathology*. **60**(1): 2–14.
3. FRAC. (2024). FRAC Code List: Fungicides sorted by mode of action. Fungicide Resistance Action Committee.
4. Garrett, K. A., Dendy, S. P., Frank, E. E., Rouse, M. N., and Travers, S. E. (2006). Climate change effects on plant disease: Genomes to ecosystems. *Annual Review of Phytopathology*. **44**(1): 489–509.
5. Harman, G. E. (2006). Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology*. **96**(2): 190–194.
6. Kulik, T., Bilska, K. and Żelechowski, M. (2020). Promising perspectives for detection, identification, and quantification of plant pathogenic fungi and oomycetes through targeting mitochondrial DNA. *International Journal of Molecular Sciences*. **21**(7): 2645.
7. Michielse, C. B., and Rep, M. (2009). Pathogen profile update: *Fusarium oxysporum*. *Molecular Plant Pathology*. **10**(3): 311–324.
8. Strange, R. N., and Scott, P. R. (2005). Plant disease: A threat to global food security. *Annual Review of Phytopathology*. **43**: 83–116.
9. Sunil, C. K., Jaidhar, C. D. and Patil, N. (2023). Systematic study on deep learning-based plant disease detection or classification. *Artificial Intelligence Review*. **56**(12): 14955-15052.
10. Wang, H., La Russa, M. and Qi, L. S. (2016). CRISPR/Cas9 in genome editing and beyond. *Annual review of biochemistry*. **85**(1): 227-264.