AGRI MAGAZINE
(International E-Magazine for Agricultural Articles)
Volume: 02, Issue: 12 (December, 2025)
Available online at http://www.agrimagazine.in
©Agri Magazine, ISSN: 3048-8656

Compatibility of Biopesticides with Other Pest Management Practices in Horticulture

*Om Narayan Nath and Tikam Das Vaishnav
RNT College of Agriculture, Kapasan (MPUAT, Udaipur), Rajasthan, India
*Corresponding Author's email: omnarayannath2007@gmail.com

Biopesticides, derived from natural sources such as microorganisms, plants, or minerals, represent a sustainable alternative to synthetic chemical pesticides in horticulture. They target pests, weeds, and diseases with minimal environmental impact, aligning with Integrated Pest Management (IPM) principles that emphasize ecological balance. However, their effectiveness largely depends on compatibility with other pest management practices, including chemical pesticides, cultural controls, biological agents, and mechanical methods.

This article discusses how biopesticides integrate into horticultural systems, highlights synergies and limitations, and provides practical insights into maximizing their effectiveness. As horticulture includes fruits, vegetables, ornamentals, and spice crops, maintaining crop health while minimizing chemical residue is essential. Compatibility ensures that biopesticides work harmoniously with other control methods without reducing efficacy or causing antagonistic effects.

Types of Biopesticides and Their Mechanisms

Understanding biopesticide categories is essential for evaluating compatibility.

1. Microbial Biopesticides

These include:

- **Bacteria** (e.g., *Bacillus thuringiensis* – Bt)
- **Fungi** (e.g., *Beauveria bassiana*, *Trichoderma harzianum*)
- **Viruses** (e.g., Baculoviruses)

They infect or suppress pests through parasitism or toxin production and often perform best under favorable environmental conditions (humidity, mild temperatures).

2. Biochemical Biopesticides

Derived from natural substances such as:

- Neem oil (azadirachtin)
- Plant extracts
- Insect pheromones

3. Plant-Incorporated Protectants (PIPs)

Genetically modified plants producing their own protective compounds (e.g., Bt crops). These are less commonly used in traditional horticulture.

The compatibility of these biopesticides depends on their mode of action, persistence, environmental sensitivity, and interaction with other pest control practices.

Integration with Integrated Pest Management (IPM)

Biopesticides play a vital role in IPM, serving as eco-friendly alternatives and enhancing system sustainability.

Synergies with Cultural Controls

Practices like crop rotation, sanitation, timely irrigation, and intercropping naturally reduce pest incidence and complement biopesticide use.

Example:

- Marigold intercropping with vegetables helps suppress nematodes, improving the performance of neem-based biopesticides.

Compatibility with Biological Controls

Many biopesticides are highly selective and safe for beneficial organisms such as ladybird beetles, parasitoids, and predatory mites.

- *Trichoderma* spp. control soil-borne pathogens without harming beneficial insects.
- Bt sprays are generally safe for pollinators when applied in the evening.

Mechanical and Physical Controls

Traps, nets, sticky boards, and pruning enhance biopesticide effectiveness. Example:

- Combining pheromone traps with microbial sprays reduces moth populations in orchards.

IPM systems often position biopesticides as the first line of defense, using chemical pesticides only when pest levels exceed economic thresholds.

Compatibility with Chemical Pesticides

Positive Interactions

Some biopesticides show synergistic effects with low-dose chemicals.

- Neem oil combined with selective insecticides reduces the required chemical dose.
- Bt mixed with pyrethroids improves control of lepidopteran pests due to complementary modes of action.

Challenges and Incompatibilities

- Chemical fungicides may kill beneficial fungi like *Beauveria* or *Trichoderma*.
- High-residue synthetic pesticides reduce survival of microbial biopesticides.
- pH-dependent degradations occur in tank mixtures.

For example:

- Copper fungicides reduce the efficacy of *Bacillus subtilis*–based biopesticides.

Management Measures

- Follow label instructions for application order.
- Apply biopesticides first and wait 2–3 days before using chemicals.
- Avoid tank-mixing unless compatibility is proven.
- Use IRAC and FRAC groups to plan rotation and avoid resistance.

Case Studies in Horticultural Crops

1. Tomato Production

Integration of *Verticillium lecanii* with sticky traps and predator release reduced whitefly infestation.

- Chemical use decreased by 60%.
- Biopesticides provided up to 80% pest control when applied early.

2. Apple Orchards

Codling moth management using granulovirus plus pheromone mating disruption resulted in reduced pesticide residues and better pollinator safety.

3. Strawberry Cultivation

Spinosad (biochemical biopesticide) combined with row covers and beneficial nematodes showed high compatibility.

- Improved soil health
- Lowered fungal disease incidence
- Increased fruit quality

These case studies show that biopesticides perform best in combination with monitoring tools like scouting, traps, and proper timing.

Challenges and Solutions

Challenges

- Sensitivity to UV radiation and temperature
- Short residual life

- Slow action compared to chemicals
- Risk of resistance if overused
- Improper mixing with incompatible pesticides

Solutions

- **Training and Awareness:** Extension programs teaching farmers about timing, mixing, and storage.
- **Improved Formulations:** Encapsulated and nano-based biopesticides enhance stability.
- **Regulatory Support:** Updated labels include compatibility guidelines.
- **Research Advances:** CRISPR-based microbial strains promise higher persistence.
- **Good Agricultural Practices (GAP):** Proper irrigation, pruning, and sanitation improve overall compatibility.

Conclusion

Biopesticides represent a cornerstone of sustainable horticulture, offering environment-friendly pest management solutions. Their compatibility with cultural, mechanical, biological, and even chemical control practices enhances their effectiveness within an IPM framework. While some challenges exist—such as environmental sensitivity and chemical incompatibility—proper training, strategic application, and advanced formulations can overcome these limitations. When used wisely, biopesticides reduce chemical dependence, safeguard beneficial organisms, improve soil health, and ensure safer produce for consumers. Their integration into horticultural pest management not only strengthens crop resilience but also contributes to long-term ecological balance.