AGRI MAGAZINE
(International E-Magazine for Agricultural Articles)
Volume: 02, Issue: 12 (December, 2025)
Available online at http://www.agrimagazine.in
©Agri Magazine, ISSN: 3048-8656

From Cure to Crisis: The Challenge of Fungicide Resistance

*Sulekha Pradhan[1], R. Arutselvan[1] and Jisha Lagna[2]
[1]ICAR-Central Tuber Crop Research Institute, Regional Station, Bhubaneswar, India
[2]Odisha University of Agriculture and Technology, Bhubaneswar, India
*Corresponding Author's email: pinkypradhan04@gmail.com

Fungicides have long stood as one of agriculture's most effective defenses against devastating fungal diseases, safeguarding global food production and shaping modern crop protection systems. However, the very success of these chemicals has fueled a crisis: the rapid rise of fungicide resistance. As fungi evolve to withstand once-reliable treatments, farmers, scientists, and public health specialists face increasing challenges in controlling both plant and human pathogens. This article explores the journey "from cure to crisis" by tracing the rise of fungicide use, explaining how resistance emerges through biological mechanisms such as target-site mutations, efflux pumps, and metabolic adaptations, and illustrating its real-world consequences through global case studies. Factors accelerating resistance ranging from overuse in agriculture to climate change and international trade are examined to highlight the interconnected pressures driving this trend. Finally, the article discusses current and emerging management strategies, including integrated pest management, fungicide rotation, resistant cultivars, and biological alternatives, offering a path forward to sustain effective disease control. Together, these insights underscore the urgent need for coordinated global action to mitigate fungicide resistance before it undermines both agricultural productivity and public health.

Key words: Fungicide resistance, target-site mutations, efflux pumps and metabolic adaptations

Introduction

For decades, fungicides have been celebrated as one of modern agriculture's greatest success stories. From protecting staple crops like wheat, rice, and potatoes to preventing spoilage in fruits and vegetables, these chemicals have played a vital role in safeguarding global food production. Without them, many of the world's most important crops would be vulnerable to destructive fungal pathogens capable of wiping out entire harvests. In an era where feeding a growing population is already a challenge, fungicides have long been viewed as an essential line of defense.

Yet this success has come with a growing and troubling cost. As fungicides have become more widely used, fungi have begun to adapt. Much like antibiotic resistance in bacteria, repeated exposure to the same fungicidal compounds has allowed fungal pathogens to evolve survival strategies, rendering once-powerful treatments increasingly ineffective. What was once a reliable cure is now turning into a crisis. Reports of fungicide-resistant strains are rising worldwide, threatening both agricultural productivity and human health.

This emerging problem is not confined to farms. Some medical antifungal drugs share chemical similarities with agricultural fungicides, meaning resistance developed in the environment can spill over into clinical settings. Pathogens such as *Aspergillus fumigatus* and *Candida auris* already difficult to treat are showing alarming levels of resistance, raising concern among public health experts.

Understanding how we arrived at this point requires looking at the evolution of fungicide use, the biological mechanisms behind resistance, and the human-driven factors

accelerating its spread. Equally important is exploring how farmers, scientists, and policymakers can respond. The challenge is clear: we must balance the need to protect crops with the need to preserve the effectiveness of fungicides for the future. This article traces the journey from cure to crisis and explores what can be done to prevent a world where fungicides no longer work.

The Rise of fungicides:

The widespread use of fungicides began in the mid-19th century, revolutionizing crop protection. Early multi-site fungicides, such as copper-based compounds, were effective but required large quantities and frequent applications. Later came "single-site" fungicides chemicals that target specific fungal processes. These modern fungicides were more potent, needed smaller doses, and offered precise modes of action. They quickly became indispensable in protecting high-value crops like wheat, grapes, potatoes, and strawberries.

Their success, however, encouraged heavy and repeated use. Global agriculture grew increasingly dependent on these chemicals, often applying them in the same fields and against the same pathogens year after year. This created ideal conditions for fungi to adapt, survive, and eventually outsmart the chemicals designed to kill them.

Fungicide Resistance and Its Mechanisms

Fungicide resistance occurs when fungal populations evolve ways to survive chemicals designed to kill them. Although the underlying genetics can be complex, the main mechanisms can be explained in simple, accessible terms:

- **Target-site mutations:** Small genetic changes alter the fungicide's binding site within the fungus, preventing the chemical from attaching and doing its job.
- **Efflux pumps:** Some fungi activate or overproduce proteins that act as molecular pumps, pushing fungicides out of their cells before they can cause damage.
- **Metabolic detoxification:** In this mechanism, fungi break down or convert fungicides into harmless compounds using specialized enzymes. This "detoxification" allows fungi to neutralize the chemical before it becomes lethal, and it can spread quickly within fungal populations when fungicides are used repeatedly.
- **Reduced uptake:** Certain fungi limit the absorption of fungicides, reducing the amount that enters the cell and weakening the chemical's effect.

These mechanisms evolve because fungal communities produce vast numbers of spores. A rare genetic change that helps one spore survive a fungicide dose can rapidly multiply, eventually leading to entire populations becoming resistant. Over time, this makes many once-effective fungicides far less reliable. (Fig.1)



Fig 1 : Key Mechanisms of Fungicide Resistance in Fungal Pathogens, (A) Reduced penetration, (B) Active efflux, (C) Metabolic detoxification, (D) Target-site modification.

Global Case Studies: Resistance in Action:

Fungicide resistance is no longer a theoretical concern; it is a lived reality around the world.

- **Downy mildew of Grapes: *Plasmopara viticola***

Plasmopara viticola, the pathogen responsible for downy mildew in grapes, has become a significant example of fungicide resistance in India. During the 2018–19 growing seasons in Karnataka, repeated use of Metalaxyl led to widespread field-level resistance across several vineyards. Resistance also emerged to QoI fungicides such as Azoxystrobin, driven by frequent and unrotated applications. Research confirmed loss of sensitivity to Metalaxyl through greenhouse trials, while a mutation in the cytochrome b gene (G143A) was identified as the key factor behind QoI resistance. Management strategies eventually succeeded through careful fungicide rotation, the use of multi-site protectants like copper oxychloride (COC) and mancozeb, and the integration of biological control agents such as *Trichoderma*, helping restore effective disease control (Massi et al.,2021).

- **Downy mildew of Fruit Crops & Vegetables: *Botrytis cinerea***

Botrytis cinerea, the gray mold fungus, is one of the most notorious examples of rapid fungicide resistance. Affecting strawberries, grapes, tomatoes, and numerous vegetables, it has repeatedly adapted to nearly every fungicide class introduced over the past five decades. Site-specific fungicides such as benzimidazoles, dicarboximides, QoIs, and SDHIs have all faced resistance soon after their release, driven by the fungus's high genetic variability and prolific spore production. Many regions now report multi-resistant Botrytis strains, particularly in strawberry fields, making disease control increasingly challenging (Hahn M., 2014). This case highlights how intensive spray programs and high disease pressure can accelerate the loss of chemical effectiveness.

Table 1: Fungicide Classes vs. Resistance Status

Fungicide Class	Mode of Action	Example Fungicides	Resistance Status	Key Pathogens
Azoles	Sterol biosynthesis inhibition	Tebuconazole, Propiconazole	Widespread	*Septoria*, *Aspergillus*
Strobilurins	QoI inhibitors	Azoxystrobin, Pyraclostrobin	High	Powdery mildew, rusts
SDHIs	Succinate dehydrogenase inhibition	Boscalid, Fluxapyroxad	Emerging	*Botrytis*, *Septoria*
Multi-site	Broad-spectrum	Mancozeb, Chlorothalonil	Low	Wide range

Factors Driving Resistance Development

Fungicide resistance does not emerge by accident—it develops when biological traits of pathogens interact with farming practices, chemical properties, and environmental pressures. Several key factors accelerate this process:

- **Pathogen Biology:** Some fungi naturally evolve resistance more quickly because of their biology. Pathogens that reproduce rapidly, produce large numbers of spores, or have high genetic diversity—like *Botrytis cinerea* or *Zymoseptoria tritici*—can generate resistance mutations faster. Airborne pathogens also spread resistant strains over long distances with ease.
- **Fungicide Characteristics:** The type of fungicide greatly influences resistance risk. Single-site fungicides, which target one specific biochemical process, are highly effective but easier for fungi to overcome through single mutations. In contrast, multi-site fungicides like mancozeb pose lower resistance risk but may be used less frequently in modern programs. The persistence, mobility, and mode of action of a fungicide all shape how quickly resistance can develop.

- **Agronomic Practices:** How crops are grown strongly impacts resistance. Monoculture farming and dense plantings create ideal conditions for diseases to build up. Overuse or repeated spraying of the same fungicide group increases selective pressure, while poor crop rotation, excessive nitrogen use, and inadequate field sanitation allow resistant strains to thrive. Inconsistent or sub lethal dosing can further speed up resistance evolution.
- **Environmental and Operational Factors:** Global movement of seedlings, fruits, and soil helps resistant spores travel across borders. Climate change brings warmer, more humid conditions that favor fungal growth and increase fungicide use. At the same time, shared chemical classes between agriculture and medicine—especially azoles—create **cross-resistance** in environmental pathogens that can impact human health. Operational factors, such as limited access to diverse fungicides or lack of technical guidance, also contribute to misuse and faster resistance buildup.

Management Strategies

Fungicide resistance is a serious threat, but smart practices can slow its spread and extend the life of these tools. Key strategies include:

- **Rotate and mix fungicides**: Alternate or combine products with different modes of action (check FRAC codes). This lowers selective pressure on fungi.
- **Apply sparingly**: Spray only when needed—scout fields and use preventive timing to reduce the number of applications.
- **Adopt integrated pest management (IPM)**: Combine fungicides with resistant crop varieties, crop rotation, biological controls, and cultural practices like removing crop debris.
- **Follow label guidelines**: Use full recommended doses and ensure thorough coverage for maximum effectiveness.

Conclusion

Fungicides have been essential allies in the fight against crop diseases, but their future effectiveness cannot be taken for granted. The rise of fungicide resistance is a warning sign that overdependence on chemical solutions is unsustainable. Yet it also presents an opportunity: to shift toward smarter, more resilient agricultural systems that balance chemical, biological, and cultural practices. Protecting the effectiveness of fungicides is not only critical for food security but also for safeguarding human health. By understanding how resistance emerges and adopting proactive strategies, we can prevent today's cure from becoming tomorrow's crisis—and ensure a healthier, more sustainable future for agriculture and beyond.

References

1. Massi, F., Torriani, S. F. F., Borghi, L., & Toffolatti, S. L. (2021). *Fungicide resistance evolution and detection in plant pathogens: Plasmopara viticola as a case study*. Microorganisms, 9(1), Article 119. https://doi.org/10.3390/microorganisms9010119.
2. Hahn, M. (2014). *The rising threat of fungicide resistance in plant pathogenic fungi: Botrytis as a case study*. Journal of Chemical Biology, 7(3), 133–141. https://doi.org/10.1007/s12154-014-0113-1.
3. Singh, R., & Kumar, P. (2022). Molecular mechanism of resistance to different fungicides: A review. International Journal of Research and Analytical Reviews, 9(2), 1570–1580.