



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 02 (February, 2026)

Available online at <http://www.agrimagazine.in>

© Agri Magazine, ISSN: 3048-8656

CRISPR/Cas-Mediated Genome Editing for Precision Plant Breeding

*Braj Mohan Singh Baghel

Senior Technical Assistant, ICAR-CSSRI, Karnal and Ph.D. Scholar, Genetics and Plant Breeding, ICAR-IARI New Delhi, India

*Corresponding Author's email: brajsbi90@gmail.com

The increasing demand for food, coupled with climate change, limited natural resources and emerging biotic and abiotic stresses, has placed unprecedented pressure on global agriculture. Conventional plant breeding approaches, although successful, are time-consuming and often imprecise due to genetic recombination and environmental influences. The advent of CRISPR/Cas-mediated genome editing has revolutionized precision plant breeding by enabling accurate, efficient and targeted modification of plant genomes. CRISPR/Cas technology allows breeders to directly edit genes associated with yield, stress tolerance, disease resistance and nutritional quality without introducing foreign DNA. This article presents a comprehensive and reader-friendly overview of the principles of CRISPR/Cas genome editing, its molecular mechanism, major applications in crop improvement, advantages over traditional breeding, challenges, regulatory considerations and future prospects. CRISPR/Cas technology represents a transformative tool for developing high-yielding, resilient and climate-smart crop varieties essential for sustainable agriculture and global food security.

Keywords: CRISPR/Cas, Genome editing, Precision plant breeding, Crop improvement, Biotechnology

Introduction

Plant breeding has played a central role in agricultural development by improving crop yield, quality and adaptability. Traditional breeding methods rely on phenotypic selection and hybridization, which require multiple generations and are often influenced by environmental factors. Although molecular markers have improved selection efficiency, the process of introducing desirable traits remains slow and limited by linkage drag. Genome editing has emerged as a breakthrough in plant biotechnology by enabling precise and predictable genetic modifications. Among genome editing tools, CRISPR/Cas systems have gained widespread attention due to their simplicity, efficiency and versatility. Since their first successful application in plants, CRISPR/Cas technologies have rapidly transformed modern plant breeding, offering unprecedented opportunities for precise genetic improvement.

Origin and Working Principle of CRISPR/Cas Technology

CRISPR/Cas technology originates from a natural defense mechanism found in bacteria and archaea, where it functions as an adaptive immune system against viral infections. Bacteria store fragments of viral DNA in CRISPR arrays and use them to recognize and destroy invading genetic material. In plant genome editing, this system is reprogrammed using a guide RNA (gRNA) designed to match a specific target DNA sequence. The gRNA directs the Cas nuclease (commonly Cas9) to the target site in the plant genome, where it introduces a precise double-strand break. The plant's endogenous DNA repair machinery then repairs the break, leading to targeted genetic modifications.




Figure 1: CRISPR/CAS Mechanism for Targeted Plant Genome Editing.

DNA Repair Mechanisms in CRISPR Editing

Non-Homologous End Joining (NHEJ)

NHEJ is the most common repair pathway in plants. It often introduces small insertions or deletions at the cleavage site, resulting in gene knockouts. This method is widely used to disable undesirable genes controlling susceptibility or negative traits.

Homology-Directed Repair (HDR)

HDR uses a homologous DNA template to introduce precise genetic changes such as gene replacement or insertion. Although less efficient in plants, HDR enables precise allele modification and trait improvement.

Types of CRISPR/Cas Systems Used in Plants

Several CRISPR/Cas variants have been adapted for plant breeding:

- Cas9: Most widely used system; efficient and versatile
- Cas12a (Cpf1): Produces staggered DNA cuts and improves editing precision
- Base Editing: Allows single-base substitutions without DNA breaks
- Prime Editing: Enables highly precise and programmable genome modifications

These advanced tools further enhance the accuracy and scope of precision breeding.

Applications of CRISPR/Cas in Plant Improvement

Abiotic Stress Tolerance

CRISPR/Cas has been used to enhance tolerance to drought, salinity and heat by editing genes involved in stress perception, ion transport, osmotic regulation and antioxidant defense mechanisms.

Disease and Pest Resistance

Genome editing enables targeted modification of susceptibility genes, resulting in improved resistance to bacterial, fungal and viral diseases. This approach provides durable resistance without relying on chemical pesticides.

Yield and Plant Architecture Improvement

Genes controlling flowering time, tillering, grain size and plant height have been edited to enhance yield potential and adaptability in major field crops.

Nutritional Quality Enhancement

CRISPR/Cas technology has been successfully applied to improve protein quality, vitamin content, oil composition and reduction of anti-nutritional factors, contributing to improved human nutrition.

Advantages of CRISPR/Cas Over Conventional Breeding

- High precision and specificity
- Rapid development of improved varieties
- Minimal or no linkage drag
- Multiplex genome editing capability
- Possibility of transgene-free edited plants

These advantages make CRISPR/Cas a powerful tool for modern precision breeding.

Challenges and Limitations

Despite its potential, CRISPR/Cas technology faces several challenges, including off-target mutations, limited efficiency of HDR, difficulties in plant transformation and regeneration and regulatory uncertainty. Continuous technological improvements are addressing these limitations.

Regulatory and Ethical Considerations

Regulatory frameworks for genome-edited crops vary across countries. Some nations regulate transgene-free edited plants similarly to conventionally bred crops, while others apply GMO regulations. Transparent, science-based policies are essential for the responsible adoption of CRISPR technology in agriculture.

Future Prospects of CRISPR-Based Precision Breeding

The integration of CRISPR/Cas with genomics, transcriptomics, artificial intelligence and high-throughput phenotyping will further accelerate crop improvement. Genome editing of orphan crops and climate-resilient varieties will play a vital role in sustainable and inclusive agricultural development.

Conclusion

CRISPR/Cas-mediated genome editing represents a paradigm shift in plant breeding by enabling precise, rapid and efficient genetic improvement. Through targeted modification of genes controlling yield, stress tolerance, disease resistance and nutritional quality, CRISPR/Cas offers powerful solutions to global agricultural challenges. Its strategic integration with conventional breeding and molecular tools will be essential for achieving sustainable food security in a changing climate.

References

1. Chen, K., Wang, Y., Zhang, R., Zhang, H., and Gao, C. (2019). CRISPR/Cas genome editing and precision plant breeding. *Annual Review of Plant Biology*, 70, 667–697.
2. Jaganathan, D., Ramasamy, K., Sellamuthu, G., Jayabalan, S., and Venkataraman, G. (2018). CRISPR for crop improvement: An update review. *Frontiers in Plant Science*, 9, 985.
3. Zhang, Y., Zhang, F., and Qi, Y. (2017). CRISPR/Cas9 genome editing in plants: Progress and challenges. *Annual Review of Plant Biology*, 68, 1–26.
4. Bortesi, L., and Fischer, R. (2015). The CRISPR/Cas9 system for plant genome editing. *Biotechnology Advances*, 33(1), 41–52.
5. Zaidi, S. S. A., Mahfouz, M. M., and Mansoor, S. (2018). Genome editing: Targeting susceptibility genes for plant disease resistance. *Trends in Biotechnology*, 36(9), 898–906.
6. Gao, C. (2021). Genome engineering for crop improvement and future agriculture. *Cell*, 184(6), 1621–1635.