



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 03 (March, 2026)

Available online at <http://www.agrimagazine.in>

© Agri Magazine, ISSN: 3048-8656

Secondary Metabolites: The Core of Plant Immunity

*Mitali Tiwari and Ishika Joshi

Ph.D. Scholar, Department of Genetics and Plant Breeding, College of Agriculture, G. B. Pant University of Agriculture & Technology, Pantnagar, Uttarakhand, India

*Corresponding Author's email: mitalitiwari111@gmail.com

Plants, being sessile organisms, have evolved sophisticated defense strategies to cope with a wide array of biotic (pathogens and herbivores) and abiotic (drought, salinity, UV radiation) stresses. Central to this defense system is the production of secondary metabolites (also known as specialized metabolites). These diverse, low-molecular-weight organic compounds including terpenoids, phenolics, and alkaloids are not directly involved in essential primary metabolic processes like photosynthesis or respiration, but they are crucial for survival. This article reviews the pivotal role of these compounds, exploring their biosynthetic pathways, their mechanisms of action as toxins, repellents, and signaling molecules, and how they collectively form the chemical core of plant immunity, offering a vast reservoir of natural defenses that can be leveraged for sustainable agriculture.

Keywords: Secondary Metabolites, Plant Immunity, Terpenoids, Phenolics, Alkaloids, Plant Defense Mechanism, Biosynthesis, Biotic Stress, Abiotic Stress

Introduction

Life on Earth is a constant struggle for survival, and for plants, this struggle is waged without the ability to flee. Over millennia, plants have developed a complex, multilayered defense system. This immunity is generally categorized into PAMP-Triggered Immunity (PTI), a broad-spectrum defense, and Effector-Triggered Immunity (ETI), a localized, strong defense. However, the *products* that ultimately execute these defense signals and protect the plant are largely the secondary metabolites. Unlike primary metabolites (e.g., carbohydrates, proteins, nucleic acids) which are essential for life, secondary metabolites are restricted in their distribution, often unique to a specific plant family or genus, and are typically produced in response to environmental cues. Their ecological roles are vast, serving as color pigments, aroma compounds, and, most importantly, chemical weapons and protective shields. The biosynthesis of these compounds requires significant metabolic investment, highlighting their critical importance to plant fitness and survival, especially under stressful conditions. Plant secondary metabolites are structurally classified into three major groups, each originating from distinct primary metabolic pathways:




Fig 1: Mechanisms of tolerance in plants and resistance to biotic and abiotic stresses

1. Terpenoids (Isoprenoids)

Source Pathway: The Methylerythritol Phosphate (MEP) pathway in the plastids and the Mevalonate (MVA) pathway in the cytosol.

- **Defense Role:** Terpenoids are derived from the basic five-carbon unit isopentenyl diphosphate (IPP). They function as volatile repellents (e.g., monoterpenes and sesquiterpenes found in essential oils), which deter generalist herbivores. Non-volatile forms, like phytoecdysteroids, can mimic insect hormones, disrupting their life cycle. A critical group are the diterpenes, which often act as phytoalexins (antimicrobial compounds produced *de novo* upon infection). For example, resin acids in conifers physically trap and chemically disable insects and microbes.

2. Phenolics

Source Pathway: The Shikimic Acid Pathway and the Malonic Acid Pathway.

- **Defense Role:** These compounds contain at least one aromatic ring with one or more hydroxyl groups. Their diversity is vast, including flavonoids, lignins, tannins, and coumarins.
- ✓ Lignins provide physical defense by strengthening the cell wall, making it an impassable barrier against microbial invasion.
- ✓ Tannins are highly effective anti-feedants due to their ability to bind and precipitate proteins, drastically reducing the digestibility of plant tissues for herbivores.
- ✓ Flavonoids (like anthocyanins) and coumarins act as UV protectants (abiotic stress defense) by absorbing harmful radiation and are potent antioxidants, neutralizing the Reactive Oxygen Species (ROS) produced during stress.

3. Alkaloids

Source Pathway: Primarily derived from amino acids (e.g., lysine, tyrosine, tryptophan).

- **Defense Role:** Alkaloids are generally cyclic compounds containing nitrogen, often possessing high levels of toxicity to animals due to their interference with neurotransmitter systems. They are the most widely known chemical toxins used by plants.
- ✓ **Nicotine** (derived from ornithine/arginine) in tobacco is a powerful neurotoxin against insects.
- ✓ **Caffeine** (a purine alkaloid) is toxic to many insects and inhibits the germination of competing plant seeds (allelopathy).
- ✓ Many alkaloids, such as those found in the poppy or nightshade families, have evolved specifically to disrupt the central nervous system of vertebrate herbivores, making them highly effective defense agents.




Fig 2: Different classes of secondary metabolites (Upadhyay *et al.*, 2025)

Signaling and Priming

Beyond their direct toxic or structural roles, secondary metabolites and their precursors act as crucial signaling molecules. For instance, the degradation products of glucosinolates (a type of defense compound found in brassicas) are volatile signals that activate defense genes in neighboring plants. The phytoalexins are, by definition, produced as a rapid signal-response mechanism, illustrating the direct link between secondary metabolism and ETI.

Table1: Secondary metabolites in the management of phytopathogens (Goura *et al.*, 2025)

Crop	Secondary metabolite	Pathogen
<i>Triticum spp.</i>	Lignin	<i>Puccinia graminis</i>
<i>Zea mays</i>	Caffeic acid	<i>Glomerella Graminicola</i>
<i>Allium sativum</i>	Alkaloids, glycosides, tannins, terpenoids, flavonoids, and saponins	<i>A. solani</i>
<i>Hyoscyamus niger</i>	Tropane alkaloids	<i>Seudomonas putida</i> , <i>P. fluorescens</i>
<i>Citrus reticulata</i>	Reticine A	TMV
<i>Nicotiana attenuata</i>	Chlorogenic acid, catechin	<i>Trichobaris mucorea</i>
<i>Oryza sativa</i>	Phytoalexin	<i>Sogatella furcifera</i> (Hemiptera: <i>Delphacidae</i>)
<i>Cajanus platycarpus</i>	Flavonoid	<i>Helicoverpa armigera</i>

Table 2: Effects on secondary metabolite production in plants under different abiotic stresses (Goura *et al.*, 2025)

Abiotic stress	Secondary Metabolite class	Plant species
UV-B	Flavonoid	<i>Arabidopsis</i> (<i>Arabidopsis thaliana</i>)
High UV radiation	Alkaloid	<i>Clematis terniflora</i>
Blue light	Anthocyanin	<i>Solanum lycopersicum</i>
Drought	Flavonoid	Wheat leaves
Salinity	Anthocyanin	<i>Arabidopsis thaliana</i>
Low temperature	Morphine	<i>Papaver somniferum</i> L.
High temperature	Ginsenoside	<i>Panax ginseng</i> C.A. Mey
Ozone	Anthocynine, Rosmarinic acid, Tannine,	<i>Melissa officinalis</i>
Heavy metal	Flavonoids	<i>Vitis vinifera</i>

Conclusion

Secondary metabolites are undeniably the chemical core of plant immunity. They provide a dynamic and versatile array of defenses, transitioning from structural fortifications (lignin) to volatile repellents (terpenes) and powerful cytotoxins (alkaloids). The strategic, yet energy-intensive, production of these compounds underscores their absolute necessity for plant survival against relentless environmental challenges. Understanding the complex biosynthetic pathways and regulatory networks governing these molecules is critical. By applying this knowledge through metabolic engineering and sustainable agricultural practices, researchers can enhance the natural chemical arsenal of crops, leading to plants that are inherently more resistant to pests, diseases, and the escalating pressures of climate change.

References

1. Akbar, R., Sun, J., Bo, Y., Khattak, W. A., Khan, A. A., Jin, C., ... & Du, D. (2024). Understanding the influence of secondary metabolites in plant invasion strategies: a comprehensive review. *Plants*, 13(22), 3162.
2. Crockett, S.L., and Wildung, M.R. (2018). Evolution of a structural diversification platform: Biosynthesis of defense compounds in the plant kingdom. *The Plant Cell*, 30(2), 268–282.
3. Dixon, R. A. (2001). Natural products and plant disease resistance. *Nature*, 411(6839), 843–847.
4. Goura, K., Legrifi, I., Kallali, N. S., Taoussi, M., Kenfaoui, J., Meddich, A., ... & Lahlali, R. (2025). Beyond survival: the role of secondary metabolites in plant defense mechanisms. *Journal of Crop Health*, 77(4), 121.
5. Isah, T. (2019). Stress and Defense Responses in Plants: The Role of Secondary Metabolites. *Biological Research*, 52(1), 39.
6. Khan, A., Kanwal, F., Ullah, S., Fahad, M., Tariq, L., Altaf, M. T., ... & Zhang, G. (2025). Plant secondary metabolites—Central regulators against abiotic and biotic stresses. *Metabolites*, 15(4), 276.
7. Kumar, S., Korra, T., Thakur, R., Arutselvan, R., Kashyap, A. S., Nehela, Y., ... & Keswani, C. (2023). Role of plant secondary metabolites in defence and transcriptional regulation in response to biotic stress. *Plant stress*, 8, 100154.
8. Kumar, S., Korra, T., Thakur, R., Arutselvan, R., Kashyap, A. S., Nehela, Y., ... & Keswani, C. (2023). Role of plant secondary metabolites in defence and transcriptional regulation in response to biotic stress. *Plant stress*, 8, 100154.
9. Piasecka, A., Jedrzejczak-Rey, N., & Bednarek, P. (2015). Secondary metabolites in plant innate immunity: conserved function of divergent chemicals. *New Phytologist*, 206(3), 948-964.
10. Pichersky, E. and Raguso, R.A. (2018). Why do plants produce so many metabolites? *Current Opinion in Plant Biology*, 48, 1–8.
11. Upadhyay, R., Saini, R., Shukla, P. K., & Tiwari, K. N. (2025). Role of secondary metabolites in plant defense mechanisms: A molecular and biotechnological insights. *Phytochemistry Reviews*, 24(1), 953-983.
12. Wink, M. (2010). Introduction: biochemistry, physiology and ecological functions of secondary metabolites. *Annual plant reviews volume 40: Biochemistry of plant secondary metabolism*, 1-19.
13. Wink, M. (2015). Modes of action of secondary metabolites from plants in defense and communication. *Current Opinion in Plant Biology*, 25, 41–48.
14. Yadav, P., & Siddiqui, Z. H. (2023). Role of Secondary Metabolites in Stress Management Under Changing Climate Conditions. *Global Climate Change and Plant Stress Management*, 383-397.
15. Zaynab, M., Fatima, M., Abbas, S., Sharif, Y., Umair, M., Zafar, M. H., & Bahadar, K. (2018). Role of secondary metabolites in plant defense against pathogens. *Microbial pathogenesis*, 124, 198-202.