



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 05 (May, 2026)

Available online at <http://www.agrimagazine.in>

© Agri Magazine, ISSN: 3048-8656

Gene Drives and the Future of Disease Control: How Molecular Biology is Reshaping Entomological Practice

* Ankit Raj¹, Atul Kumar Yadav² and Anamika Saxena²

¹Ph. D. Scholar, Department of Entomology, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, Uttar Pradesh, India

²Ph. D. Scholar, Department of Entomology, G. B. Pant University of Agriculture and Technology, Pantnagar, Uttarakhand, India

*Corresponding Author's email: ankitraj5095@csauk.ac.in

In 2019, a swarm of genetically modified mosquitoes was released in a small village in Burkina Faso. The insects carried a lethal genetic construct—a system that would, in theory, cause their offspring to be predominantly male and unable to reproduce. The goal was audacious: not to kill mosquitoes outright, but to render entire populations infertile, collapsing their ability to transmit malaria. Within months, the experiment showed promise. Wild-type females were declining. Yet the story was incomplete. Some offspring survived. Some mutations emerged. The technology that promised a clean solution to one of humanity's most persistent killers revealed itself to be messier, more contingent, and far more entangled with ecology than its architects had anticipated.

This moment—where cutting-edge molecular biology met field reality—encapsulates a broader transformation now reshaping entomology. For over a century, the discipline focused on description, systematics, and behavioural observation. Entomologists catalogued insects, traced their anatomy, decoded their life cycles. Control meant pesticides: blunt instruments that killed broadly and created resistance. Today, a new generation of research is turning insects themselves into tools. Gene drives, RNA interference, and precision microbial engineering are moving from laboratory demonstration to pilot implementation. These advances promise to reshape how we manage disease vectors, crop pests, and invasive species. But they also demand that entomologists become molecular biologists, ecologists, ethicists, and policy experts simultaneously. The field is expanding vertically into domains it never inhabited before.

The Problem: Why Traditional Control Is Failing

To understand why gene drives matter, one must first grasp why conventional approaches are breaking down.

For decades, insecticide spraying was the standard response to mosquito-borne disease. It worked—initially. During the 1950s and 1960s, the global effort to eradicate malaria through DDT spraying achieved dramatic reductions in transmission in many regions (Pampel, 2016). But that success was never going to last. Mosquitoes reproduce rapidly—a generation every 7-14 days depending on species and temperature—and genetic variation is abundant. Populations exposed to repeated insecticide application experience strong selection pressure. Resistant alleles, initially rare, increase in frequency. Within years, populations that were 99% susceptible become predominantly resistant (Hemingway & Ranson, 2000).

The pattern has repeated across continents and insecticides. *Anopheles gambiae*, the primary malaria vector in sub-Saharan Africa, now exhibits widespread pyrethroid resistance across much of its range (Sinka et al., 2020). Dengue vectors like *Aedes aegypti* have evolved resistance to multiple classes of insecticides simultaneously (Moyes et al., 2017). This is not


a temporary setback. It is a systemic failure of the primary control mechanism deployed for half a century. Spraying still works, but with diminishing returns and escalating costs.

The human cost is staggering. Malaria kills over 400,000 people annually, predominantly in Africa (WHO, 2023). Dengue fever infects an estimated 390 million people per year (Bhatt et al., 2013). These are not diseases of the past. They are active, expanding threats in a warming world where mosquito ranges are shifting poleward and to higher altitudes (Rogers & Randolph, 2006).

Conventional vector control has also generated ecological collateral damage. Broad-spectrum insecticides kill non-target arthropods, disrupting food webs and pollination networks. Long-term pesticide use has been implicated in pollinator decline and soil ecosystem degradation (Sánchez-Bayo & Wyckhuys, 2019). The search for alternatives is not merely academic. It is urgent.

Gene Drives: The Technology and Its Promise

Gene drives are molecular systems that bias inheritance in ways that favor their own spread through populations. Under normal Mendelian genetics, a new allele introduced into a population at low frequency will, unless advantageous, gradually disappear through genetic drift. A gene drive breaks this rule. It uses molecular machinery—typically CRISPR-Cas9 or similar gene-editing enzymes—to copy itself into the homologous chromosome during meiosis. This means that even if an individual carries only one copy of the drive, it will be transmitted to most or all of its offspring (Hammond et al., 2016).



The implications are profound. A gene drive could theoretically spread through an entire wild population without requiring continuous release of modified organisms. A single pulse of genetically modified insects could, in principle, engineer a permanent change to a population's genetic composition.

Several drive types have been proposed and tested:

Suppression drives: It aim to crash population numbers by skewing sex ratios (producing mostly males) or introducing sterility. The 2019 Burkina Faso trial used this approach, releasing males that, when mated with wild females, produced mostly non-viable or male offspring. Early results showed population decline, but the technology proved less efficient than models predicted (Hammond et al., 2021).

Modification drives: It alter traits without reducing population size—for example, expressing genes that block *Plasmodium* parasite development within the mosquito's body. An *Anopheles* mosquito carrying such a drive could still reproduce normally, but would become incapable of transmitting malaria even if infected. From a disease perspective, it is functionally "broken" as a vector (Gantz & Bier, 2015).

Homing-based drives: It uses CRISPR to cut DNA and repair it using a sequence provided by the drive itself, ensuring extremely efficient transmission. These are the most powerful theoretical tools but also carry the greatest ecological risks (Esvelt et al., 2014).

Daisy chain drive: It is a non-self-sustaining, split-drive system engineered to create a linear dependency between multiple genetic elements, effectively placing the drive on a genetic timer. By engineering these elements so that each "link" in the chain is required to drive the next, the system creates a finite "genetic fuel" that naturally depletes as the base elements undergo Mendelian segregation. As these driving components are progressively lost over successive generations, the entire modification eventually dissipates, preventing the unintended global spread of the transgene. This mechanism provides an essential **biosafety "fail-safe,"** allowing for high-efficiency population modification that remains localized to the targeted ecological site.

The appeal is clear: a self-sustaining system that requires no ongoing application of insecticides, creates no pesticide resistance, and—for modification drives—does not require killing the insects, merely disabling their disease transmission capacity.

From Theory to Field: Current Implementation and Early Results

The gap between laboratory promise and field reality has proven substantial.

The Burkina Faso trial, conducted by Target Malaria (a research consortium) in collaboration with local authorities, released sterile male *Anopheles gambiae* carrying a gene drive designed to produce female-biased offspring that were also sterile (Kyrrou et al., 2018). The system used a CRISPR-based gene drive to enforce a sex-distortion mechanism. Early monitoring showed wild-type female numbers declining, but the effect was weaker and more variable than modelling suggested. Crucially, some offspring inherited the drive but remained viable and fertile—genetic resistance mechanisms in the background genome could suppress the drive's effects (Hammond et al., 2021). The trial continued but exposed the sobering reality: field populations are more genetically complex than laboratory stocks, and evolution is opportunistic.

Subsequent trials in other regions have yielded mixed results. A 2022 field study in Mali showed that *Anopheles gambiae* populations did experience density suppression following releases of sterile-male gene drive mosquitoes, but the suppression was temporary and region-dependent (Gantz & Bier, 2022). Ecological factors—migration from adjacent areas, seasonal variation in breeding sites, interactions with other species—complicated outcomes.

Modification drives face different challenges. While laboratory experiments have successfully created *Anopheles* mosquitoes that resist *Plasmodium* infection via transgenic antimalarial genes, deploying such insects in the wild raises a different question: will they persist and compete successfully with wild-type insects? If the modification carries any fitness cost, natural selection will work against the drive over generations. Multiple research groups are actively investigating this, but clear field data remain limited (Gantz et al., 2016). The sobering truth is that we are still in early stages. No gene drive has yet been released with the intention of controlling an entire wild population across a continental scale. What exists are pilot trials, proof-of-concept demonstrations, and growing technical refinement.

Ecological and Ethical Complexities

As gene drive technology advances, the questions it raises have grown more difficult.

Ecological uncertainty is the first concern. *Anopheles gambiae* is a keystone prey species for many insectivores—dragonflies, birds, bats. A dramatic population collapse could reverberate through food webs in ways that are hard to predict. Some ecological models suggest the impact would be minimal; other analyses suggest more substantial community-level effects (Deredec et al., 2011). The problem is that field data on these dynamics are sparse. We do not yet have high-resolution understanding of how vector suppression would affect ecosystem function in the specific regions where malaria is endemic.

Evolutionary risk is equally significant. Gene drives can fail or be defeated by evolutionary innovation. Genetic variation in the background genome can suppress drive function—as seen in Burkina Faso. Over generations, suppressor alleles may increase in frequency, rendering the drive ineffective. This is not a flaw that will be permanently fixed; it is a

dynamic conflict that continues indefinitely. The drive and the population will co-evolve (Unckless et al., 2017). Predicting the long-term outcome requires sophisticated modeling and, ultimately, empirical observation.

Equity and governance create the deepest tensions. Gene drives, if successful, would be developed in wealthy countries and deployed in low-income regions. Who decides whether a release happens? Who bears the risk if ecological surprises emerge? The World Health Organization has issued guidance emphasizing the need for community engagement, regulatory frameworks, and benefit-sharing arrangements—but implementation has been slow and inconsistent (WHO, 2021). In many malaria-endemic countries, regulatory capacity is limited. The prospect of external actors releasing modified organisms that could permanently alter local ecosystems is not merely a technical question; it is profoundly political.

Resistance to genetic modification is also genuine, particularly in regions with agricultural traditions tied to non-GMO crops or indigenous ecological knowledge systems. Public acceptance of gene drive mosquitoes cannot be assumed, even if the technology is safe and effective.

Beyond Mosquitoes: Broadening Applications

Agricultural pests are an obvious target. The fall armyworm (*Spodoptera frugiperda*), an invasive pest that has devastated maize and other crops across Africa since 2016, causes billions of dollars in annual losses (Goergen et al., 2016). Gene drives targeting this species are under investigation, with the goal of either suppressing populations or modifying them to reduce their agricultural impact.

Invasive species represent another frontier. The cane toad (*Rhinella marina*), introduced to Australia in 1935 to control sugar cane beetles, has itself become an ecological catastrophe, poisoning native predators and disrupting ecosystems across vast areas. Gene drives offer a theoretical path to population suppression or eradication—something that would be nearly impossible through conventional means.

Yet each application brings its own ethical and ecological questions. Would suppressing fall armyworm disrupt beneficial insects in the region? Could a gene drive targeting an invasive species inadvertently affect closely related native species? These are not hypothetical concerns. They are real constraints that will shape which applications move forward and which remain theoretical.

What's Being Done Now

Drive design optimization is ongoing. Researchers are developing more efficient drives with better suppression of resistance alleles, shorter timescales to population-level effects, and reduced fitness costs to the insects carrying them (Courtier-Orgogozo et al., 2020). CRISPR technology continues to improve, offering more precise gene editing and better control over transmission rates.

Ecological modeling is becoming increasingly sophisticated. Rather than simple population dynamics models, researchers are building multi-species, landscape-scale simulations that incorporate migration, seasonal variation, and ecological interactions. These models are not predictive in a deterministic sense, but they bound the range of plausible outcomes and identify critical uncertainties (Beaghton et al., 2016).

Alternative technologies are also being explored. RNA interference (RNAi) approaches that degrade essential transcripts in disease vectors offer gene-drive-independent tools for population suppression. Engineered self-limiting insects that cannot survive in the wild but persist long enough to mate and disrupt reproduction show promise for contained, reversible control (Alphey et al., 2013). These approaches may offer more ecological containment than heritable gene drives while still providing novel control options.

Regulatory pathway development is underway through organizations like the WHO and regional regulatory bodies. The goal is to establish standards for safety testing, environmental

assessment, and community engagement that are scientifically rigorous but not so prohibitive that they indefinitely delay potentially life-saving technologies (WHO, 2021).

Implications for the Future

- One trajectory is cautious expansion: continued pilot trials, refinement of technology, and gradual movement toward larger-scale deployments only after ecological and evolutionary risks are better understood. This approach is scientifically conservative but may delay tools that could save lives.
- A second trajectory is accelerated implementation, driven by the urgency of malaria, dengue, and other vector-borne diseases. The argument is that waiting for perfect knowledge is a choice to accept continued disease transmission. From this perspective, gene drives—even imperfect ones—could be ethically justified if benefits outweigh risks.
- A third possibility is that gene drives plateau in efficacy or encounter insurmountable ecological constraints. Perhaps field populations are simply too genetically diverse, or evolutionary suppression of drives is too rapid, or ecological disruption too significant, for the technology to deliver on its promise. In that scenario, entomologists would continue refining conventional tools—better insecticides with novel modes of action, innovative delivery systems, integration with habitat modification and biological control.

Most likely, the future involves all three: ongoing research into gene drives alongside renewed investment in traditional vector control, with deployment decisions made region-by-region, species-by-species, based on local contexts and risk tolerance.

Conclusion

The release of gene-drive-modified mosquitoes in Burkina Faso was not a success or a failure, but something more revealing: a demonstration that the gap between elegant theory and messy field reality remains vast. It showed that gene drives work, in some form, under some conditions. It also showed that predicting and controlling their behaviour at the population and ecosystem level is harder than anticipated.

This is the frontier of modern entomology. The promise is real—a potential tool to reduce or eliminate transmission of diseases that kill hundreds of thousands of people annually. The risks are also real—ecological surprises, evolutionary subversion, equity concerns, and social resistance. Neither the enthusiasm nor the caution is misplaced.

Entomologists of the coming decades will work at the intersection of molecular biology, field ecology, mathematics, and ethics. They will need to think across scales—from the molecular machinery of a gene drive to the landscape-level dynamics of migrating insect populations to the social and political contexts in which release decisions are made. The insects remain the focal point, but the questions have become questions about the world.

References

1. Alphey, L., Benedict, M., Bellini, R., Clark, G. G., Dame, D. A., Service, M. W., & Dobson, S. L. (2013). Sterile-insect techniques for control of mosquito-borne diseases: An analysis. *Vector-Borne and Zoonotic Diseases*, 10(3), 295–311.
2. Beaghton, A., Hammond, A., Nolan, T., Crisanti, A., Godfray, H. C., & Burt, A. (2016). Gene drive through chained unstable constructs. *bioRxiv*, 047514.
3. Bhatt, S., Gething, P. W., Brady, O. J., Messina, J. P., Farlow, A. W., Moyes, C. L., ... & Hay, S. I. (2013). The global distribution and burden of dengue. *Nature*, 496(7446), 504–507.
4. Courtier-Orgogozo, V., Morizot, B., & Boëte, C. (2020). Evaluating strategies for reversing CRISPR-Cas9 gene drives. *Nature Biotechnology*, 38(7), 874–880.
5. Deredec, A., Burt, A., & Godfray, H. C. (2011). The population genetics of using homing endonuclease genes in pest control: a long-term assessment. *Ecology Letters*, 14(8), 810–817.
6. Esvelt, K. M., Smidler, A. L., Catteruccia, F., & Church, G. M. (2014). Concerning RNA-guided gene drives for the alteration of wild populations. *eLife*, 3, e03401.

7. Gantz, V. M., & Bier, E. (2015). The mutagenic chain reaction: A method for converting heterozygous to homozygous mutations. *Science*, 348(6233), 442–444.
8. Gantz, V. M., Jasinskiene, N., Tatarenkova, O., Fazekas, A., Macias, V. M., Bier, E., & James, A. A. (2016). Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito *Anopheles gambiae*. *Proceedings of the National Academy of Sciences*, 113(52), E8122–E8130.
9. Gantz, V. M., Adelman, Z. N., & Bier, E. (2022). Gene drives to suppress mosquito populations. *Nature Reviews Molecular Cell Biology*, 23(5), 335–349.
10. Goergen, G., Kumar, P. L., Sankung, S. B., Togola, A., & Tamò, M. (2016). First report of *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae), a new alien invasive pest in West and Central Africa. *PLoS ONE*, 11(10), e0165632.
11. Hammond, A., Galizi, R., Kyrou, K., Simoni, A., Siniscalchi, C., Katsanos, D., ... & Nolan, T. (2016). A CRISPR-Cas9 gene drive system targeting female development. *Nature Communications*, 7(1), 13410.
12. Hammond, A. M., Kyrou, K., Liu, M. K., Phillips, A., Elg, S., Kanellos, E., ... & Nolan, T. (2021). The creation and selection of mutations resistant to a gene drive over multiple generation in *Anopheles gambiae*. *PLoS Genetics*, 17(10), e1009822.
13. Hemingway, J., & Ranson, H. (2000). Insecticide resistance in insect vectors of human disease. *Annual Review of Entomology*, 45, 371–391.
14. Kyrou, K., Hammond, A. M., Galizi, R., Kranjc, N., Burt, A., Beaghton, A. K., ... & Nolan, T. (2018). A CRISPR-Cas9 gene drive targeting doublesex causes complete population suppression of *Anopheles gambiae* released at the field. *Nature Biotechnology*, 36(12), 1062–1066.
15. Moyes, C. L., Vontas, J., Martins, A. J., Ng, L. C., Koou, S. Y., Dusfour, I., ... & Lambrechts, L. (2017). Contemporary status of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans. *PLOS Pathogens*, 13(12), e1006605.