



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 06 (June, 2026)

Available online at <http://www.agrimagazine.in>

© Agri Magazine, ISSN: 3048-8656

Paddy Straw to Table - Growing Paddy Straw Mushroom (*Volvariella volvacea*) — A Complete Grower's Guide

*Sujitha Sri R and Dr. K. Vignesh

Palar Agricultural College, Melpatti, Vellore, Tamil Nadu, India

*Corresponding Author's email: sujiramki28@gmail.com

Paddy Straw Mushroom (*Volvariella volvacea*) mushroom, scientifically known as *Volvariella volvacea*, is a tropical edible mushroom widely cultivated in Southeast Asia due to its fast growth and high nutritional value. It grows naturally on decomposing paddy straw and is recognized for its short cropping cycle of 10–14 days, making it highly economical for farmers. The mushroom is rich in Proteins, vitamins B & C, minerals like potassium and phosphorus, and contains low fat and calories, making it ideal for health-conscious diets.

Cultivation requires high temperature of 28–35°C and 80–90% relative humidity, which suits tropical and subtropical climates. Substrates like paddy straw, cotton waste, and banana leaves are commonly used after pasteurization. Key steps in cultivation include spawn preparation, bed preparation, spawning, and maintaining proper moisture and ventilation.

Major challenges include contamination by molds, short shelf life, and sensitivity to temperature fluctuations. Despite this, its medicinal properties such as antioxidant, anti-inflammatory, and cholesterol-lowering effects increase its demand. Value addition through drying, pickling, and powdering helps improve shelf life and marketability. The mushroom also aids in agricultural waste management by converting paddy straw into protein-rich food. Recent studies focus on improving yield through better strains and controlled environment farming. Government schemes and training programs are promoting its cultivation for rural employment and income generation. In conclusion, paddy straw mushroom is a sustainable, nutritious, and profitable crop with strong potential in food security and agribusiness. Future research should target extending shelf life and developing disease-resistant varieties for large-scale commercial production.

A Mushroom Born from the Tropics

Walk through any bustling market in Southeast Asia or coastal India and you are likely to spot a cluster of small, dome-shaped mushrooms nestled inside straw baskets their pale caps smooth and slightly velvety, their aroma faintly earthy and inviting. These are paddy straw mushrooms, known to science as *Volvariella volvacea*, and their story is one of remarkable resourcefulness. They thrive on the same rice straw that farmers would otherwise burn at field edges, converting agricultural waste into protein-rich, flavourful food within just three weeks. This article takes you through every stage of paddy straw mushroom cultivation from understanding the biology of the fungus to harvesting marketable fruiting bodies in language accessible to farmers, students, and curious food enthusiasts alike. Why Grow *Volvariella volvacea*? Nutrition and Economics

Nutritional Powerhouse in a Petite Package

On a dry-weight basis, *Volvariella volvacea* contains roughly 20 to 30 percent protein comparable to legumes along with a generous complement of essential amino acids that are often limited in plant-based diets. It is rich in B-complex vitamins (thiamine, riboflavin, niacin), vitamin C, and dietary fibre, while being naturally low in fat and calories. Its mineral

content potassium, phosphorus, calcium, and iron among them makes it nutritionally attractive to communities where micronutrient deficiency is a concern.



Economic Returns

The crop's economics are equally appealing. A standard bed prepared from 10 kg of dry paddy straw can yield between 1.0 and 1.5 kg of fresh mushrooms within 20 to 25 days. Fresh paddy straw mushrooms command premium prices in urban markets and hotels. Beyond fresh sales, drying, pickling, and powder-making extend shelf life dramatically and open access to wholesale and export channels.

Understanding the Growing Environment

Every successful crop begins with a thorough understanding of what the organism needs from its surroundings. *Volvarela volvacea* is a thermophilic fungus, meaning it genuinely prefers warmth a characteristic that distinguishes it sharply from temperate species like oyster mushroom or button mushroom.

Environmental Factor	Optimum Range	Effect if Out of Range
Temperature	30 – 35 °C	Below 25°C slows growth; above 38°C kills mycelium
Relative Humidity	80 – 90%	Below 70% causes pinhead abortion; above 95% promotes rot
Substrate pH	6.5 – 7.0	Acidic or alkaline conditions inhibit colonisation
Light	Diffused / low intensity	Darkness prolongs incubation; strong direct light is harmful
Ventilation	Moderate air exchange	Poor ventilation causes CO ₂ build-up and distorted fruiting

Getting Started: Materials and Substrate Selection

What You Will Need

The beauty of paddy straw mushroom cultivation lies in the simplicity of its requirements. Most materials are either freely available on farms or inexpensive to procure:

Dry, clean paddy straw the primary raw material

- High-quality mushroom spawn freshly prepared or sourced from a reliable supplier
- Clean water for soaking and daily moisture maintenance
- Polythene sheets or fresh banana leaves for covering beds during incubation
- Rope or twine to bundle and hold straw layers
- Sanitation supplies bleach, lime, or wood ash for site disinfection

Choosing the Right Substrate

Paddy straw is the benchmark substrate, prized for its high cellulose content, appropriate particle size, and widespread availability in rice-growing regions. When selecting straw, look for bundles that are golden-yellow in colour, odour-free, and completely free of mould. Straw that has been stored outdoors and is partially decomposed or discoloured by black or green patches should be rejected it harbours competing organisms that will overwhelm the mushroom mycelium. Farmers in areas where paddy straw is scarce have successfully cultivated the species on several alternative substrates, each with its own advantages and limitations:

Banana leaves and pseudostem: readily decomposable, high moisture retention, excellent for coastal and humid regions

Cotton gin waste: rich in cellulose, produces high yields but benefits from pasteurisation

- Sugarcane trash and bagasse: widely available in cane-growing zones; works best when mixed with straw
- Water hyacinth: an invasive aquatic weed converted into a useful input a win for both grower and environment

Substrate Preparation: Setting the Stage for Growth

Soaking

Dry straw is bundled into manageable portions and fully submerged in a tank, drum, or lined pit containing clean water. The straw absorbs water over a period of 12 to 18 hours. This rehydration phase is critical: the mycelium requires a substrate moisture content of 65 to 70 percent to colonise efficiently. An easy field test squeeze a handful of soaked straw firmly; only a few drops should fall, not a stream of water. After soaking, the bundles are lifted out and stacked at an angle on a raised surface for 30 to 60 minutes to allow excess water to drain away.

Pasteurisation (Optional but Recommended)

Raw straw carries dormant bacteria, fungal spores, and insect eggs that can outcompete the mushroom mycelium during the early colonisation phase. Pasteurisation heating the substrate to 60 – 80°C for one hour using steam or hot-water immersion dramatically reduces this microbial competition without sterilising the substrate completely. A lightly pasteurised substrate retains some beneficial microorganisms that help suppress pathogens later in the cropping cycle. Growers who skip pasteurisation often achieve adequate results in clean, ventilated environments during cooler periods, but the technique is strongly recommended for beginners and for cultivation during the rainy season when contamination pressure is highest.

Site Sanitation

The growing area whether an open yard, a shade structure, or a dedicated mushroom house should be cleaned and disinfected before bed construction begins. Sweeping away organic debris and washing the floor and raised platform surfaces with a dilute bleach solution (1%) or lime wash prevents carry-over contamination from previous crops.

Bed Construction and Spawning

Building the Bed

Beds are built on raised platforms bamboo racks, wooden tables, or cemented slabs elevated 30 to 50 cm above the ground to facilitate drainage and reduce pest pressure from ground-dwelling insects. A convenient bed size is approximately 1 metre wide by 1 metre long, which allows easy access for watering and harvesting from all sides. Construction proceeds

layer by layer. Spread a 10 cm base of soaked straw on the platform and press it lightly to create an even, compact surface. Sprinkle spawn along a 5 cm strip around all four edges and lightly across the central area. Add a second straw layer, press gently, and repeat the spawning pattern. Continue building in this manner until three or four layers are complete, finishing with a thin covering layer of straw over the topmost spawn.

Spawn Rate and Quality

Spawn is applied at approximately 5 to 7 percent of the dry weight of the straw used. For a bed prepared from 10 kg of dry straw, this translates to 500 to 700 g of spawn usually about one standard spawn packet. Using more spawn than this does not proportionally increase yield and is wasteful; using too little results in slow colonisation and higher contamination risk. It should be covered in dense white mycelium, smell pleasantly of mushrooms (not sour or ammonia-like), and show no discoloration.

Covering the Bed

Once construction is complete, the bed is draped with a polythene sheet or overlapping banana leaves to trap warmth and moisture during the initial colonisation phase. Leave the base of the bed slightly open or ensure that the covering is not sealed airtight a small amount of gas exchange helps the mycelium breathe without allowing the surface to dry out.

Incubation, Pinning, and Harvest

The Incubation Phase

After the bed is covered and left undisturbed, the mycelium begins its invisible work. Over the following 10 to 15 days, white fungal threads ramify through the straw, bonding the fibres into a dense, interconnected mass. During this period, the bed temperature should be maintained between 30 and 35°C. Avoid opening the cover unnecessarily each exposure allows warm air to escape and cold dry air to enter, slowing colonisation. A fully colonised bed feels firm and springy, and the straw is visibly bound together by white mycelial growth.

Inducing Fruiting

Once colonisation is complete, remove the cover fully or fold it back to allow light and fresh air to reach the bed surface. Mist the surface lightly with clean water. The change in temperature, humidity, and carbon dioxide concentration that follows signals the mycelium to shift from vegetative growth to reproductive mode. Within two to three days, tiny white pinheads will emerge the first visible signs of the fruiting bodies that will develop into harvestable mushrooms.

Water Management During Fruiting

Consistent surface moisture is essential during pinning and fruiting. Mist the bed two or three times daily using a fine-spray nozzle, ensuring the straw surface stays moist but not waterlogged. Water droplets sitting in the cup-shaped volva at the base of developing mushrooms can encourage bacterial rot. Always mist with a gentle, upward-angled spray that wets the straw rather than saturating the fruit bodies directly.

Harvesting at the Egg Stage

The paddy straw mushroom must be harvested before the outer membrane the universal veil ruptures and the cap expands. Once the veil tears, the mushroom releases spores, its shelf life collapses within hours, and its market value drops sharply. The ideal harvest stage is the egg stage: the fruiting body is oval to round, the veil is smooth and intact, and the overall diameter is between 3 and 5 cm. The mushroom will remain marketable for 24 to 48 hours at ambient tropical temperatures. To harvest, grasp the base of the mushroom firmly and give a gentle twist-and-pull motion. Avoid using a knife if possible, as cutting leaves a wound in the substrate that can become a point of bacterial entry. After harvesting each cluster, remove any remnants of torn veil or damaged tissue from the bed surface.

Post-Harvest Handling

Volvariella volvacea is one of the most perishable of all cultivated mushrooms. Its high water content (around 90%) and vigorous cellular respiration mean that, at room temperature, quality deteriorates within 12 to 24 hours of harvest.

Immediate Post-Harvest Care

- Handle harvested mushrooms as little as possible to prevent bruising
- Pack loosely in ventilated bamboo baskets or perforated polythene bags
- Keep in the coolest available location and transport to market within hours of harvest
- If refrigeration is available, store at 10 – 15°C (not lower temperatures below 8°C damage the tissue)

Protecting the Crop: Diseases and Pests

Even a well-managed bed is not immune to problems. Recognising symptoms early and responding promptly is the key to minimising losses.

Problem	Symptoms	Cause	Management
Green mould (Trichoderma spp.)	Green or blue-green powdery patches on substrate	Contaminated spawn or poor pasteurisation	Remove affected portion; improve sanitation
Bacterial soft rot	Slimy, foul-smelling fruiting bodies	Excess moisture + warm temperature	Reduce watering; improve air circulation
Cobweb mould	Fine grey webbing over bed surface	High CO ₂ and stagnant air	Increase ventilation; light lime dusting
Insect damage (flies, mites)	Holes in fruit bodies; stunted pins	Poor sanitation, nearby organic waste	Neem spray; keep surroundings clean

Prevention is always more effective than treatment. A discipline of thorough site sanitation, careful spawn handling, and attentive moisture management eliminates the vast majority of contamination problems before they begin.

The Road Ahead: Scope and Opportunity

India's climate and agricultural geography place it in an excellent position to significantly expand paddy straw mushroom production. States with year-round warm temperatures Tamil Nadu, Andhra Pradesh, Telangana, Odisha, West Bengal, and Kerala among them offer natural growing conditions that require minimal supplemental inputs. Growing consumer awareness of the health value of mushrooms, the expansion of organised retail, and the rising popularity of plant-based diets are collectively driving demand upward.

Conclusion

The paddy straw mushroom is a gift from the tropics fast-growing, nutritious, economically accessible, and environmentally responsible. By mastering a handful of straight forward techniques, any motivated grower can transform a pile of rice straw into a marketable, nutritious crop in under a month. Supplement farm income, or build a scalable mushroom enterprise, *Volvariella volvacea* offers a compelling starting point and the techniques described in this guide will take you confidently from the first soaked straw bundle to the final, satisfying harvest.

References

1. Indian Council of Agricultural Research. (1988). Mushroom Cultivation. New Delhi: ICAR.
2. Pathak, V. N., Yadav, N., & Gaur, M. (2000). Mushroom Production and Processing Technology. Jodhpur: Agrobios Publishers.
3. Stamets, P. (2000). Growing Gourmet and Medicinal Mushrooms (3rd ed.). Berkeley: Ten Speed Press.
4. Tripathi, D. P. (2005). Mushroom Cultivation. Jaipur: Oxford Book Company.
5. Singh, R. P., & Chaube, H. S. (1996). Mushroom Cultivation and Processing. Jodhpur: Scientific Publishers.

6. Harini M, Reddy PSK, Vignesh K, Hari K, Reddy PSS. Unlocking the genetic diversity of bush hyacinth bean (*Lablab purpureus* var. *typicus* L.) landraces under South Indian agro-ecologies. Genet Resour Crop Evol. 2025;72:1-24.
7. Vignesh K. Green synthesis and characterization of zinc oxide nanoparticles using neem (*Azadirachta indica* L.) leaf extract. Asian J Plant Pathol 2025; 19(1): 27–35. <https://doi.org/10.1016/j.mtcomm.2023.105942>
8. Vignesh K. Survey on the incidence of Fusarium wilt of tomato incited by *Fusarium oxysporum* f. sp. *lycopersici* (Fol) in major tomato growing areas of krishnagiri district. Plant Archives. 2021;21:2369-2376
9. K. Vignesh, L. Vengadeskumar, S. Sanjaygandhi, T. Sabesan, Prevalence of maydis leaf blight of maize in Tamil Nadu and assess the morphological character and virulence of *Bipolaris maydis* (NISIK.) shoemaker, The Pharma Innovation Journal, 12, 6, 65-70 (2023)
10. Yadav, M. C., Verma, R. N., & Bahl, N. (2012). A Text Book on Mushroom Production Technology. Solan: Directorate of Mushroom Research, ICAR.
11. Chang, S. T., & Miles, P. G. (2004). Mushrooms: Cultivation, Nutritional Value, Medicinal Effect, and Environmental Impact (2nd ed.). Boca Raton: CRC Press.