



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 06 (June, 2026)

Available online at <http://www.agrimagazine.in>

© Agri Magazine, ISSN: 3048-8656

Cut Foliage Fillers and Their Importance

Amrutha S S¹, *Shakila Sadasivam², Sona V¹ and Susindhar Babu E¹

¹PG Scholar, Department of Floriculture and Landscaping, SRM College of Agricultural Sciences, Baburayanpettai, Chengalpattu -603201, Tamil Nadu

²Asst. Prof., Department of Floriculture and Landscaping, SRM College of Agricultural Sciences, Baburayanpettai, Chengalpattu -603201, Tamil Nadu

*Corresponding Author's email: shakilas@srmist.edu.in

Cut foliage fillers also known as the cut greens which are leaves and leafy stems used in floral arrangements, bouquets and interior decorations (Mirunalini *et al.*). These cut foliage fillers play a vibrant role in enhancing floral decorations by adding colour, contrast and volume lastly structure of the end products like arrangements and bouquets and gives a complete look. A study shows that about 25% to 30% of bouquets include foliage, that highlights the importance of foliage fillers that are used in floriculture industry (Mirunalini *et al.*). Thus, cut foliage fillers are not only attractive but also an essential component in the floriculture industry which combines aesthetic beauty and have high economic value.

Characteristics of Cut Foliage Fillers

- The foliage should have long vase life
- The foliage fillers should be in attractive colour and texture
- Good keeping quality
- Transport stress resistance

Uses of Cut Foliage Fillers








- Used in Bouquets and floral arrangements
- Wedding and event decor
- Interior decoration and indoor scaping
- Dry flower arrangements



Economic importance




Cut foliage fillers represents a significance in the floriculture industry due to their low production cost and high market demand. They are particularly suitable for small scale farmers as they require less investment compared to the flower production. There is strong demand for foliage in international markets such as Europe, USA and Japan. Florida has been a leading producer of cut foliage and contributes to the floriculture economy (Jhanji *et al.*).

Environmental and Health Importance

Cut foliage plants play an important role in promoting green living and human health. Many foliage species contribute to improving indoor air quality by removing harmful pollutants like carbon dioxide, formaldehyde, benzene and nitrogen oxides. Prolonged exposure to these pollutants may lead to respiratory diseases, fatigue and some other long term health complications or issues. In addition to that foliage plants help maintain optimal indoor humidity levels, which reduces dryness in the air and enhances overall well-being in indoor spaces (Talaie and Ponraj, 2024). The presence of indoor greenery has also help to lower stress levels improved emotional well-being and better functioning of everyday life these makes the foliage plants highly valuable in green buildings, workplaces and healthcare centers. (Wang *et al.*, 2023).

Common Name	Scientific name	Uses	
Amaranthus	<i>Amaranthus tricolor</i>	Used as colourful foliage in bouquets and floral arrangements. Available in red, green and yellow colours.	
Aglaonema	<i>Aglaonema commutatum</i>	Used as decorative foliage filler due to its variegated leaves.	
Alpinia	<i>Alpinia purpurata</i>	Used in large arrangements and background filling as it have bold large foliage.	
Croton	<i>Codianum variegatum</i>	Popular for its bright, multi coloured foliage. Used to create texture in floral arrangements.	
Dracaena	<i>Dracaena marginata</i>	Used for its structural foliage. It provides height in floral arrangements.	
Peace lily	<i>Spathiphyllum wallisii</i>	Used in indoor décor and in floral arrangements. Both leaves and flowers are used for its graceful appearance.	
Aspidistra	<i>Aspidistra elatior</i>	It is known for its strong and long-lasting leaves. Widely used in floral art, wrapping foliage and ikebana	
Arrowhead plant	<i>Syngonium Podophyllum</i>	Used as a soft foliage filler to add delicate texture and volume in bouquets.	

Baby's Breath	<i>Gypsophylla paniculata</i>	Adds a soft effect in bouquets. Used in wedding bouquets and decorations.	
Golden rod	<i>Solidago canadensis</i>	Used to create natural look in arrangements. Enhances warmth in the floral decoration with its golden yellow colour.	
Statice	<i>Limonium sinuatum</i>	Provides papery texture and vibrant colour in the floral arrangements and bouquets.	
Wax flower	<i>Chamelaucium uncinatum</i>	Used in bridal bouquets and corsages. Blends well with both pastel and bold coloured flowers.	
Thuja	<i>Thuja orientalis</i>	Widely used as a filler and background material in bouquets. Provides a soft, feathery texture and bright green colour.	
Asparagus fern	<i>Asparagus setaceus</i>	Ideal for filling empty spaces in floral arrangements. Fine, feathery foliage gives the fuller view.	
Sword fern	<i>Nephrolepis exaltata</i>	Used in bouquets and interior floral decoration. Ideal for softening the appearances of arrangements.	
Leatherleaf fern	<i>Rumohra adiantiformis</i>	Widely used as a base and background filler in bouquets. Provides fullness and uniform green texture.	

Tree fern	<i>Asparagus virgatus</i>	Used to provide volume and structural support and suitable for large-scale floral decorations and stage décor.	
Monstera	<i>Monstera deliciosa</i>	Popularly used in modern and exotic floral designs. Large, split leaves used for focal arrangements.	
Areca palm	<i>Dypsis lutescens</i>	Popularly used for its foliage as background for a fuller look.	

References

1. Mirunalini, S. P., Shanthi, A., Manikandan, M., Jayalakshmi, C., Ganga, M., Bhargavi, M. S. N. V. S. S. P., & Sumera, S. (2026). *Cut Foliage Crops: Production Systems, Postharvest Handling and Global Trade Perspectives*. Journal of Scientific Research and Reports, 32(4), 107–125.
2. Jhanji, S., Kaur, G., & Singh, P. (2026). Enhancing Postharvest Quality of Cut Foliage Through Active Modified Atmosphere Packaging. New Zealand Journal of Crop and Horticultural Science, 54(1), e70026.
3. Wang, Y., Zhang, Z., Xu, Y., Rodgers, T. F., Ablimit, M., Li, J., & Tan, F. (2023). Identifying the contributions of root and foliage gaseous/particle uptakes to indoor plants for phthalates, OPFRs and PAHs. Science of The Total Environment, 883, 163644.
4. Talaie, A., and Ponraj, M. (2024). Impact of houseplants on reducing indoor air pollution. Journal of Environmental Treatment Techniques, 12(3), 1- 13.