



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 06 (June, 2026)

Available online at <http://www.agrimagazine.in>

© Agri Magazine, ISSN: 3048-8656

Effect of Climate Change on Insect Pest Population

*Anjali Swarnkar and Manoj Kumar Mahla

Department of Entomology, Rajasthan College of Agriculture, MPUAT,
Udaipur, Rajasthan, India

*Corresponding Author's email: swarnkaranjali40@gmail.com

Climate change has emerged as one of the most significant challenges to global agriculture and food security. Increasing concentrations of greenhouse gases resulting from industrialization, urbanization, deforestation, and intensive agricultural activities have led to rising global temperatures, altered precipitation patterns, and more frequent extreme weather events. These changes have profound effects on agricultural ecosystems, particularly on insect pests, which are highly sensitive to environmental conditions due to their poikilothermic nature. Climatic factors such as temperature, humidity, rainfall, and atmospheric carbon dioxide (CO₂) directly influence insect growth, development, reproduction, survival, migration, and geographical distribution. Rising temperatures can accelerate insect metabolism, increase voltinism, reduce overwintering mortality, and facilitate the expansion of pest species into previously unsuitable regions. In addition, elevated CO₂ concentrations and changing climatic conditions can modify host plant physiology and disrupt pest–natural enemy interactions, thereby increasing the likelihood of pest outbreaks. In India, the emergence and spread of economically important pests such as fall armyworm (*Spodoptera frugiperda*), rice planthopper (*Nilaparvata lugens*), pink bollworm (*Pectinophora gossypiella*), and sugarcane woolly aphid (*Ceratovacuna lanigera*) have highlighted the growing impact of climate change on crop protection. Therefore, understanding the effects of climate change on insect pest populations is essential for developing sustainable and climate-resilient pest management strategies.

Effects of Rising Temperature on Insect Pest Populations

Temperature is considered the most important climatic factor regulating insect biology and population dynamics. Most insect species possess an optimum temperature range within which growth, development, survival, and reproduction occur most efficiently. An increase in temperature within this favourable range accelerates physiological processes and shortens developmental duration, leading to rapid population growth. One of the most significant consequences of climate warming is the increase in voltinism, or the number of generations completed per year. Shorter developmental periods enable many pest species to complete additional generations during a cropping season, thereby increasing their population density and potential for crop damage. Higher temperatures also enhance reproductive rates in many insects, resulting in greater fecundity and faster population build-up. Climate warming influences insect survival as well. Reduced winter mortality allows larger populations to survive adverse seasons and initiate infestations earlier in the following crop season. Earlier emergence and prolonged activity periods increase feeding duration and crop exposure to pest attack. Furthermore, rising temperatures may facilitate the establishment of invasive insect species in new regions where climatic conditions were previously unfavourable. Temperature also affects insect migration and geographical distribution. Numerous studies have documented poleward and upward shifts in insect species ranges in response to warming trends. Such changes may introduce new pests into agricultural systems and increase the vulnerability of crops to emerging pest threats.

Effects of Elevated Atmospheric Carbon Dioxide

Atmospheric carbon dioxide concentration has increased substantially since the industrial revolution and is projected to continue rising in the future. Elevated CO₂ affects insect pests primarily through changes in host plant physiology and nutritional quality. Increased CO₂ often enhances plant growth but reduces nitrogen concentration in plant tissues, resulting in lower nutritional quality for herbivorous insects. To compensate for reduced nitrogen availability, many chewing insects increase their feeding rates, causing greater damage to crop plants. Elevated CO₂ may also alter plant defensive compounds, influencing insect growth, development, and reproductive performance. In some cases, aphid populations have shown increased reproduction under elevated CO₂ conditions, whereas other insects exhibit reduced growth rates or altered developmental patterns. The interaction between elevated CO₂ and insect pests is complex and species-specific. Therefore, understanding these interactions remains essential for predicting future pest problems under changing climatic conditions.

Effects of Rainfall and Humidity on Insect Pests

Rainfall and relative humidity are important climatic factors that influence the abundance, survival, distribution, and outbreak potential of insect pests. Changes in precipitation patterns resulting from climate change can directly affect insect populations or indirectly influence them through changes in host plant growth and microclimatic conditions. High humidity generally favors the development and reproduction of several sap-feeding insects, whereas excessive rainfall may reduce pest populations through direct mortality, displacement, or disruption of feeding activities. Altered rainfall patterns can create favourable conditions for pest outbreaks in many cropping systems. Armyworm outbreaks have frequently been associated with periods of heavy rainfall and flooding. Similarly, increased humidity and moderate temperatures have been reported to favor the population build-up of *Maruca vitrata* in pigeon pea and several sucking pests in vegetable crops. In contrast, heavy rainfall may suppress aphid populations by washing individuals from plant surfaces and reducing their feeding efficiency. Climate-induced changes in rainfall distribution are therefore expected to influence pest incidence, seasonal abundance, and geographical spread. Drought conditions also affect insect pest populations. Water-stressed plants often undergo physiological changes that may alter their susceptibility to herbivorous insects. Some insect species benefit from drought stress due to reduced plant defensive responses, whereas others experience reduced survival and reproductive success. Consequently, changing precipitation regimes can significantly alter pest–host plant interactions and agroecosystem stability.

Effects of Climate Change on Insect Distribution and Migration

Climate change has emerged as a major driver of changes in insect distribution and migration patterns. Rising temperatures enable many insect species to expand their geographical ranges into regions that were previously unsuitable for their survival and reproduction. Such distributional shifts are commonly observed toward higher latitudes and elevations where climatic conditions are becoming increasingly favourable. Several studies have documented climate-induced range expansion in insect populations. Warming temperatures have facilitated the establishment of agricultural pests in new regions, increasing the risk of crop damage and economic losses. Changes in climatic conditions can also influence migration behavior by altering wind patterns, seasonal temperatures, and host plant availability. Migratory insects often respond rapidly to climatic fluctuations, leading to changes in migration timing, distance, and frequency. Climate change also affects insect phenology, resulting in earlier emergence, altered seasonal activity, and extended periods of pest infestation. Earlier spring emergence has been reported in several aphid and lepidopteran species, increasing their synchronization with susceptible crop growth stages. These phenological changes can significantly enhance pest pressure and challenge existing pest management strategies.

Climate Change and Insect Pest Outbreaks

One of the most important consequences of climate change is the increased frequency and severity of insect pest outbreaks. Rising temperatures, altered rainfall patterns, and elevated atmospheric CO₂ concentrations create favourable conditions for rapid pest multiplication and survival. Increased voltinism, reduced winter mortality, and expansion into new habitats contribute significantly to outbreak development. In India, several major pest outbreaks have been associated with changing climatic conditions. The rapid spread of fall armyworm (*Spodoptera frugiperda*) across maize-growing regions is considered one of the most significant examples of climate-linked pest expansion. Similarly, outbreaks of rice planthopper (*Nilaparvata lugens*) have caused severe yield losses in rice ecosystems under favorable temperature and humidity conditions. Pink bollworm (*Pectinophora gossypiella*) has re-emerged as a major threat to cotton production, while sugarcane woolly aphid (*Ceratovacuna lanigera*) and papaya mealybug (*Paracoccus marginatus*) have demonstrated remarkable adaptability to changing environmental conditions. Climate change is also increasing the likelihood of invasive insect species establishing in new agricultural regions. Such invasions often result in severe economic losses because natural enemies and effective management strategies may be absent in newly colonized areas. Therefore, understanding the relationship between climate variability and pest outbreaks is essential for developing early warning systems and effective pest management programs.

Impact of Climate Change on Natural Enemies

Natural enemies, including predators, parasitoids, and entomopathogens, play a crucial role in regulating insect pest populations within agroecosystems. Climate change can influence the abundance, diversity, and effectiveness of these beneficial organisms. Since natural enemies often differ from pests in their response to environmental conditions, changes in temperature and humidity may disrupt ecological balance and biological control processes. Increased temperatures may enhance predator and parasitoid activity up to an optimum level, beyond which their efficiency declines. Climate-induced changes in developmental rates can create asynchrony between pests and natural enemies, reducing the effectiveness of biological control. For example, if pests emerge earlier than their parasitoids due to warming temperatures, pest populations may escape natural regulation and increase rapidly. Changes in precipitation and humidity can also influence the survival and infectivity of entomopathogenic fungi and other microbial control agents. Consequently, climate change may alter the overall effectiveness of biological control programs and require modifications in integrated pest management strategies.

Climate-Smart Pest Management Strategies

The growing impact of climate change on insect pests necessitates the development of climate-smart pest management approaches. Traditional pest management practices may become less effective under changing environmental conditions, requiring adaptive and sustainable strategies. Integrated Pest Management (IPM) remains the most effective approach for addressing climate-related pest challenges. Climate-smart IPM emphasizes regular pest monitoring, ecological engineering, biological control, conservation of natural enemies, host plant resistance, and need-based pesticide application. Weather-based pest forecasting models can provide early warning of pest outbreaks and support timely management decisions. Advances in remote sensing, geographic information systems (GIS), artificial intelligence, and digital agriculture offer new opportunities for pest surveillance and forecasting. The development of climate-resilient crop varieties and strengthening biological control programs will further contribute to sustainable pest management under future climatic scenarios.

Future Perspectives

Future research should focus on understanding species-specific responses of insect pests to climate change and developing predictive models for pest risk assessment. Long-term

monitoring programs are essential for detecting shifts in pest distribution, phenology, and outbreak patterns. Greater attention should also be given to studying climate-induced changes in host plant resistance, pest-natural enemy interactions, and invasive pest dynamics. The integration of climate science, entomology, and digital technologies will play a critical role in strengthening agricultural resilience. Development of decision-support systems, climate-informed pest forecasting tools, and adaptive management strategies will be necessary to minimize future crop losses and ensure sustainable food production.

Conclusion

Climate change is profoundly influencing insect pest populations by altering their biology, survival, reproduction, distribution, migration, and outbreak potential. Rising temperatures, elevated atmospheric carbon dioxide concentrations, changing rainfall patterns, and extreme weather events are creating increasingly favourable conditions for many economically important insect pests. These changes pose significant challenges to agricultural productivity and food security worldwide. At the same time, climate change can disrupt natural enemy populations and reduce the effectiveness of biological control. Therefore, climate-resilient pest management strategies based on Integrated Pest Management, pest forecasting, biological control, and climate-smart agricultural practices are essential for sustainable crop protection. A comprehensive understanding of climate-pest interactions will be crucial for developing adaptive strategies capable of meeting future agricultural challenges under a changing climate.

References

1. Rehman, S. A. and Kumar, R. 2018. Scenario of insect pests under changing climatic situations. *International Journal of Chemical Studies*, **6**(3): 77-81.
2. Curnutte, L. B., Simmons, A. M. and Abd-Rabou, S. 2014. Climate change and *Bemisia tabaci* (Hemiptera: Aleyrodidae): impacts of temperature and carbon dioxide on life history. *Annals of the Entomological Society of America*, **107**(5): 933-943.
3. Subedi, B., Poudel, A. and Aryal, S. 2023. The impact of climate change on insect pest biology and ecology: Implications for pest management strategies, crop production, and food security. *Journal of Agriculture and Food Research*, **14**, 100733.
4. Skendžić, S., Zovko, M., Živković, I. P., Lešić, V. and Lemić, D. 2021. The impact of climate change on agricultural insect pests. *Insects*, **12**(5): 440.
5. Dhawan, A. K., Singh, K., Aneja, A. and Saini, S. 2009. Distribution of mealybug, *Phenacoccus solenopsis* Tinsley in cotton with relation to weather factors in South-Western districts of Punjab. *Journal of Entomological Research*, **33**(1): 59-63.