



Biochar and Carbon Neutrality: Turning Agricultural Waste into a Climate Solution

*Nilanjan Sarkar¹ and Adrija Bhandari²

¹PG Scholar, Department of Agronomy, Dr RPCAU, Samastipur, Bihar, India

²PG Scholar, Department of Agronomy, OUAT, Bhubaneswar, Odisha, India

*Corresponding Author's email: nilanjansarkar96@gmail.com

Every harvest season, millions of tonnes of crop residue go up in smoke across the world's grain belts. Stubble burning clears fields quickly and cheaply, but it also releases carbon dioxide, black carbon and particulate matter into an atmosphere already carrying more CO₂ than at any point in human history. At the same time, decades of intensive cultivation have thinned the organic carbon reserves of many agricultural soils, weakening their structure and fertility. Biochar sits at the intersection of these two problems. Rather than burning residues in the open or leaving them to decompose quickly, biochar production converts them into a stable, carbon-rich material that can be returned to the soil — turning a disposal problem into a potential climate and soil-health asset.

What Is Biochar?

Biochar is the solid, carbon-rich product formed when biomass — crop residues, wood waste, manure, or other organic feedstocks — is heated under low-oxygen conditions in a process called pyrolysis. Because oxygen is limited, the biomass does not combust fully; instead it breaks down into three co-products: biochar (solid), bio-oil (liquid) and syngas (gas). Pyrolysis reorganises the biomass carbon into fused aromatic ring structures that are far more resistant to microbial decomposition than the original plant tissue, which is the key property that distinguishes biochar from ordinary organic matter or ash.

Biochar and the Carbon Cycle

In an unmanaged system, carbon fixed by photosynthesis returns to the atmosphere within years to decades as residues decompose or burn. Pyrolysis interrupts this cycle: converting residues to biochar and applying it to soil can retain a large share of the original carbon in a form with residence times of decades to centuries or more, rather than the much shorter turnover of raw biomass (Lehmann et al., 2021).

Biochar and Carbon Neutrality

It is tempting to treat biochar as automatically “carbon-neutral” or “carbon-negative,” but this is not guaranteed. Whether a given biochar system removes net carbon from the atmosphere depends on the balance between the carbon it stores and the emissions generated in producing, transporting and applying it — not on the existence of biochar alone.

Life-Cycle Carbon Balance

A credible carbon accounting exercise must trace the full chain: feedstock growth and collection, transport, drying, pyrolysis energy demand, biochar transport and field application, the fraction of carbon retained in the biochar, any offsets from bio-oil or syngas used as energy, and the long-term persistence of the stored carbon. Only once these are netted against each other can a system be described as carbon-negative, and the outcome varies considerably by feedstock and technology pathway.

Biochar and Soil Carbon Sequestration

Once in soil, biochar's persistence depends on its chemical structure and its interactions with the surrounding environment. Highly condensed, low-hydrogen biochars tend to resist decomposition better than lightly charred material, and biochar carbon that becomes associated with soil minerals or protected within aggregates can persist even longer. Feedstock type, pyrolysis temperature, soil type, climate and application rate all influence how much carbon is retained and for how long (Lehmann et al., 2021). Recent work also suggests biochar-amended soils can additionally sorb atmospheric CO₂ through mineral-driven carbonation reactions, adding a secondary sequestration pathway alongside the stored biochar carbon itself (Gui et al., 2025).

Agronomic Benefits

Biochar can influence soil organic carbon, aggregation, bulk density, porosity, water-holding capacity, cation exchange capacity, nutrient retention, pH and microbial habitat. In principle these changes can improve nutrient-use efficiency and crop productivity, but the responses reported in the literature are highly context-dependent: outcomes shift with soil type, climate, crop and biochar properties, and biochar should not be treated as a universal substitute for fertilizer.

Biochar and Greenhouse Gases

Beyond CO₂ storage, biochar can influence soil emissions of methane and nitrous oxide. A meta-analysis of biochar's effect on soil N₂O emissions found an average reduction of roughly half, with biochars of low molar H:C_{org} ratio — indicating a high degree of aromatic condensation — proving more effective than lightly charred material, and with substantially smaller reductions typically observed under field conditions than in controlled laboratory incubations (Cayuela et al., 2015). Methane responses are less consistent, varying with soil redox conditions, moisture and land use. These effects are not universal and depend strongly on soil type, nitrogen management and cropping system.

Biochar and Circular Agriculture

Converting residues that would otherwise be burned or left to decompose into biochar, then returning that biochar to the soil, fits within a circular bioeconomy model: it recycles a farm-generated resource, reduces the need for open burning, and supports climate-smart agricultural systems without requiring new land.

Challenges and Limitations

Significant obstacles remain. Feedstock supply must be sustainable and not compete with other residue uses such as fodder or mulch. Production and transport carry their own emissions. Biochar quality varies widely with feedstock and pyrolysis conditions, and some feedstocks can introduce contaminants such as heavy metals or polycyclic aromatic hydrocarbons. Economic feasibility, uncertainty in carbon accounting methods, and inconsistent field responses across environments all complicate large-scale deployment, underscoring the need for standardized measurement and reporting methodologies.

Future Prospects

Advances in pyrolysis technology, clearer carbon-market accounting frameworks, more rigorous life-cycle assessments, and region- and soil-specific biochar recommendations could all improve the reliability of biochar's climate contribution. Independent global assessments now converge on a gigaton-scale sustainable mitigation potential for biochar systems, though estimates still vary depending on assumptions about feedstock availability and technology pathways (Weng & Cowie, 2025; Ayaz et al., 2025; Deng et al., 2024). Combining biochar with compost, integrating it into climate-smart cropping systems, and sustaining long-term field monitoring will be central to realising this potential responsibly.

Conclusion

Biochar offers a genuine opportunity to convert agricultural residues from a waste-management burden into a resource for soil improvement and long-term carbon storage. Realising that potential, however, is not automatic: it depends on the complete life cycle of the biochar system — from feedstock sourcing through pyrolysis to field application — being managed and measured with scientific rigour. Used carefully, biochar can be one credible piece of the broader climate-smart agriculture toolkit; used carelessly, it risks becoming another overstated climate fix.

References

1. Ayaz, M., Muntaha, S. T., Baltrėnaitė-Gedienė, E., & Kriauciūnienė, Z. (2025). Biochar and carbon-negative technologies: Exploring opportunities for climate change mitigation. *Biochar*, 7(1), 17. <https://doi.org/10.1007/s42773-024-00421-3>
2. Cayuela, M. L., Jeffery, S., & van Zwieten, L. (2015). The molar H:Corg ratio of biochar is a key factor in mitigating N₂O emissions from soil. *Agriculture, Ecosystems & Environment*, 202, 135–138. <https://doi.org/10.1016/j.agee.2014.12.015>
3. Deng, X., Teng, F., Chen, M., et al. (2024). Exploring negative emission potential of biochar to achieve carbon neutrality goal in China. *Nature Communications*, 15, 1085. <https://doi.org/10.1038/s41467-024-45314-y>
4. Gui, X., Xu, X., Zhang, Z., et al. (2025). Biochar-amended soil can further sorb atmospheric CO₂ for more carbon sequestration. *Communications Earth & Environment*, 6, 5. <https://doi.org/10.1038/s43247-024-01985-5>
5. Lehmann, J., Cowie, A., Masiello, C. A., Kammann, C., Woolf, D., Amonette, J. E., Cayuela, M. L., Camps-Arbestain, M., & Whitman, T. (2021). Biochar in climate change mitigation. *Nature Geoscience*, 14, 883–892. <https://doi.org/10.1038/s41561-021-00852-8>
6. Weng, Z. H., & Cowie, A. L. (2025). Estimates vary but credible evidence points to gigaton-scale climate change mitigation potential of biochar. *Communications Earth & Environment*, 6, 259. <https://doi.org/10.1038/s43247-025-02228-x>