



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 10 (October, 2026)

Available online at <http://www.agrimagazine.in>

©Agri Magazine, ISSN: 3048-8656

Harvesting Hope: How Climate-Smart Farming is Saving Bengal's Saline Coast

*Surja Barik, Aaheli Pal, Sreya Mandal and Esha Bag

Division of Agronomy, School of Agriculture and Rural Development, Ramakrishna Mission Vivekananda Educational & Research Institute, Narendrapur Campus, Kolkata, West Bengal - 700103, India

*Corresponding Author's email: surjabarik04@gmail.com

The coastal saline zone of West Bengal, including the iconic Sundarbans, stands on the frontlines of climate change. For local farmers, creeping soil salinity, seawater intrusion, violent cyclones, and increasingly erratic rainfall are daily threats to their survival. These harsh realities have trapped rural communities in a highly vulnerable cycle: relying heavily on a single monsoon-fed rice crop, while leaving fields largely barren during the dry winter season due to severe freshwater scarcity. But a powerful lifeline is emerging through Climate-Smart Agriculture (CSA). Rather than fighting a losing battle against nature, CSA equips farmers to adapt and thrive. It involves outsmarting the elements by adopting modern salt-tolerant seed varieties, actively harvesting rainwater, and using innovative “land shaping” techniques to improve drainage and store freshwater. Crucially, CSA breaks the traditional “rice-only” habit. By introducing short-duration pulses, oilseeds, and winter vegetables, farmers can transform saline fallow lands into year-round productive assets. Furthermore, integrating these diverse crops with backyard fisheries, livestock, and agroforestry builds a resilient safety net, ensuring stable incomes even when extreme weather strikes. Digital tools like localized weather advisories also put critical decision-making data directly into growers' hands. However, making this resilient vision a widespread reality requires overcoming high initial costs, fragmented landholdings, and limited extension services. By delivering affordable, location-specific interventions and strong institutional support, Bengal's vulnerable coastal shores can be transformed into a sustainable engine of rural prosperity and lasting hope.

Keywords: Climate-Smart Agriculture, Coastal saline soil, Sundarbans, Crop diversification, Climate resilience

Introduction

Global climate change is rapidly reshaping agricultural landscapes, but nowhere is this ecological shift felt more severely than in the coastal saline zone of West Bengal. Anchored by the fragile and dynamic ecosystem of the Sundarbans, this region stands on the frontlines of a climate crisis. Here, shifting temperatures and erratic rainfall are compounded by extreme localized threats: tidal flooding, violent cyclones, seawater intrusion, and rapidly escalating soil salinity. For generations, farmers in this deltaic region have walked an ecological tightrope. The traditional farming model is heavily dependent on a single Kharif (monsoon) rice crop, grown when rainwater is abundant enough to dilute soil salts. However, once the monsoon retreats, severe freshwater scarcity and salt-crusts soils make winter (Rabi) cultivation nearly impossible. This monoculture trap results in low cropping intensity, leaving farm incomes stagnant and communities highly vulnerable to the slightest weather anomaly.

To secure the future of coastal agriculture, traditional farming must evolve. Enter Climate-Smart Agriculture (CSA) a scientifically grounded, multi-dimensional approach designed to boost productivity while actively buffering against climate shocks. By blending modern agricultural science with ecological management, CSA offers a comprehensive toolkit: from breeding salt-tolerant crop varieties and deploying precision water management, to shifting toward robust Integrated Farming Systems (IFS). This article explores the unique agricultural bottlenecks strangling West Bengal's coastal zones and outlines how targeted, climate-smart strategies can transform these vulnerable landscapes into sustainable, year-round engines of food security and livelihood resilience.

The Geography of Risk and the Accelerating Climate Threat

The Ganges delta, encompassing the Sundarbans and its adjoining coastal plains, is a landscape defined by extremes. The region is characterized by low-lying, poorly drained alluvial soils that face a constant, dual threat: devastating waterlogging during the monsoon and extreme freshwater scarcity during the dry months. The greatest invisible enemy here is salt, which infiltrates the land from multiple fronts: tidal surges, creeping seawater intrusion, and saline groundwater that rises to the surface through capillary action when the topsoil dries. While the heavy monsoon rains temporarily wash the salt away, the dry winter season pulls it back up, turning once-fertile fields into toxic, salt-crusting wastelands. Today, this already delicate ecological balance is buckling under the accelerating weight of climate change. Rising sea levels and the increasing frequency of severe cyclones are driving highly destructive storm surges deeper inland, inundating freshwater ponds and ruining soil fertility for years. Simultaneously, erratic rainfall patterns are upending traditional farming calendars. Sudden, heavy downpours cause immediate flooding, while prolonged dry spells exhaust the already limited freshwater reserves, making it nearly impossible for seeds to germinate or crops to survive. Confronted by these escalating compounding threats, breaking the traditional farming mold is no longer just an option for coastal farmers; it is an absolute necessity for survival.

Breaking the Cycle: The Role of Climate-Smart Agriculture

Historically, coastal farmers have adapted to this harsh environment by relying entirely on the monsoon. They plant a single crop of Kharif rice when rainwater is abundant, leaving the land entirely fallow during the dry winter. The widespread use of long-duration rice varieties worsens this monoculture trap. Because these traditional crops take so long to mature, they delay the harvest, leaving farmers with zero time or residual soil moisture to plant a second winter (Rabi) crop. Climate-Smart Agriculture (CSA) fundamentally disrupts this cycle of vulnerability. By shifting to medium-duration rice varieties, farmers can harvest earlier, unlocking a critical window to plant a second crop while the soil still holds moisture. Coupled with climate-smart agronomic practices, farmers can then introduce pulses, oilseeds, and vegetables suited to the remaining soil conditions. This not only maximizes land use and breaks the dependency on a single monsoon harvest, but it also creates multiple streams of income, ensuring that a single extreme weather event does not wipe out a family's entire livelihood.

The Blueprint for Resilience: Core Climate-Smart Strategies

To survive and thrive in the coastal saline zone, farming can no longer rely on guesswork or outdated traditions. Climate-Smart Agriculture (CSA) deploys a combination of biological interventions, hydrological engineering, and soil rehabilitation to build a resilient farming system. Here are the core scientific strategies driving this transformation:

1. Biological Defense: Deploying Climate-Ready Genetics The first line of defense against a hostile climate is genetic. Traditional rice varieties are highly susceptible to sudden flooding and salt stress. By transitioning to scientifically bred, climate-resilient cultivars, farmers can drastically reduce their risk. For example, submergence-tolerant rice varieties like *Swarna-Sub1* and *CR Dhan 801* can survive underwater for days during cyclone-induced floods,

while *Sahbhagi Dhan* is optimized for drought-prone spells. Beyond rice, adopting stress-tolerant pulses (such as *Pusa Vishal* mungbean) allows farmers to capitalize on the limited soil moisture available during the dry winter (Rabi) season, securing a second harvest where conventional crops would fail.

2. Hydrological Engineering: Mastering Water and Topography In the Sundarbans, managing water means managing salt. The scientific solution lies in Land Shaping—mechanically altering the farm's topography into a calculated system of raised beds, ridges, furrows, and farm ponds. This serves a dual purpose: it harvests precious monsoon rainwater for winter irrigation, and it elevates the root systems of sensitive crops above the waterlogged, saline soil surface.

Furthermore, harvested freshwater is utilized for a critical process known as leaching. By applying good-quality freshwater to the fields through efficient micro-irrigation (drip or sprinkler systems), farmers can physically "wash" excess accumulated salts downward, pushing them safely below the active root zone of the crops.

3. Precision Diagnostics and Soil Rehabilitation: A cornerstone of CSA is regular, precision soil and water testing. By monitoring Electrical Conductivity (EC) and soil pH, alongside testing irrigation water for Total Dissolved Solids (TDS) and Sodium Adsorption Ratio (SAR), farmers can accurately map the toxicity of their fields and apply targeted soil amendments.

To rehabilitate degraded soils, CSA heavily promotes the use of organic matter (farmyard manure, compost, and crop residues). Organic matter acts as a biological sponge, improving soil structure, enhancing water-holding capacity, and buffering roots against salt damage.

4. The Science of Mulching: One of the most cost-effective and scientifically elegant climate-smart practices is mulching. Covering the soil surface with crop residues does more than just conserve moisture. In coastal zones, intense sun evaporates surface water, which triggers a "capillary action" that pulls highly saline groundwater up to the surface, leaving a toxic salt crust. A thick layer of mulch physically blocks this evaporation, breaking the capillary rise and keeping the root zone hospitable for plant growth.

5. Ecosystem Synergy through Integrated Farming: Ultimately, the goal of CSA is to eliminate the risks of a single-crop failure. Integrated Farming Systems (IFS) weave crop cultivation, livestock, fisheries, and agroforestry into a single, closed-loop ecosystem. In this model, farm waste is endlessly recycled: livestock manure becomes organic fertilizer, crop residues serve as animal feed, and farm ponds provide both fish and vital irrigation water. This ecological synergy drastically lowers the cost of external fertilizers and creates a diversified economic safety net that is highly resilient to climate shocks.

Strategic Crop Diversification: Breaking the Monoculture Trap

The traditional "rice-fallow" system, where fields remain completely barren post-monsoon due to salt accumulation and water stress, is a primary driver of economic instability in the region. The climate-smart alternative is strategic crop diversification. By transitioning away from a singular reliance on Kharif rice, farmers can deploy salt-resistant crops, optimize residual soil moisture, and utilize harvested rainwater to keep their lands productive year-round.

Choosing the right crop rotation is highly dependent on the specific micro-environment of the farm. Below is a breakdown of successful, location-specific cropping systems currently transforming the coastal belt:

Table 1: Climate-Smart Crop Diversification Strategies in Coastal West Bengal

Cropping System	Ideal Location / Condition	Current Status & Agronomic Impact
Rice–Mustard–Jute	Hasnabad (North 24 Parganas)	Widely documented; maximizes year-round land utilization and provides diverse income streams.

Cropping System	Ideal Location / Condition	Current Status & Agronomic Impact
Rice–Pulses	Basanti & less-saline pockets	Highly effective; utilizes residual winter moisture for hardy pulses like khesari, restoring soil nitrogen.
Rice–Vegetables	Areas with freshwater access	High-income generator; winter cultivation of chilli, potato, brinjal, and gourds using precision irrigation.
Rice–Boro Rice	Gosaba	Practiced selectively where harvested pond water is sufficient to support a second rice cycle.
Rice–Fish / Aquaculture	Low-lying Sundarban fields	Established synergy; aquatic environments provide organic nutrition to the rice while yielding a high-value protein crop.
Rice–Cotton	Highly saline rice-fallows	An emerging success (spearheaded by ICAR-ATARI, Nimpith); short-duration cotton is proving highly profitable on otherwise barren salt tracts.

The Closed-Loop Economy: Integrated Farming Systems (IFS)

To truly build climate resilience, farms must mimic natural ecosystems. Integrated Farming Systems (IFS) combine multiple enterprises such as crops, dairy, poultry, fisheries, and agroforestry into a single, self-sustaining loop where the waste of one component becomes the fuel for another. For instance, a Crop + Livestock + Fishery model drastically reduces a farmer's reliance on expensive external inputs. Crop residues feed the cattle, cattle dung generates biogas and vermicompost, and poultry droppings fertilize the fish pond. The pond, in turn, provides life-saving irrigation water for winter crops. If a sudden storm damages the rice harvest, the farmer is not left destitute, as the livestock and fisheries provide a robust economic safety net.




Fig 1: Integrated farming system Model for Costal Saline zone of West Bengal

Digital Agriculture: Empowering the Modern Farmer

Building livelihood resilience is no longer just about physical labor; it is about data. Digital and precision agriculture are taking the guesswork out of coastal farming. Today, Geographic Information Systems (GIS) and remote sensing are being utilized to map real-time soil salinity and monitor crop health from above. On the ground, affordable soil-moisture sensors and mobile agro-meteorological advisories deliver hyper-local weather forecasts directly to farmers' smartphones. This allows them to precisely time their planting, apply exact fertilizer doses, and schedule micro-irrigation only when necessary, conserving every drop of precious freshwater while protecting their yields from sudden climatic shifts.

Roadblocks and the Path Forward

Despite the clear benefits, scaling Climate-Smart Agriculture across West Bengal's coastal zone faces real-world hurdles. Farmers are constrained by highly fragmented landholdings, the steep initial costs of earth-moving for land shaping, and a lack of access to quality seeds and advanced micro-irrigation tech. Furthermore, weak extension services often leave rural farmers disconnected from the latest scientific breakthroughs. Future agricultural research and policy must bridge this gap. The focus must remain heavily on breeding "multi-stress tolerant" crops (capable of surviving floods, heat, and salt simultaneously), designing ultra-low-cost rainwater harvesting structures, and leveraging AI-based localized weather advisories.

Conclusion

The coastal saline zone of West Bengal is at a critical ecological crossroads. The compounding threats of rising soil salinity, severe freshwater scarcity, and devastating cyclones have made traditional farming methods economically and environmentally unsustainable. However, Climate-Smart Agriculture offers a proven, science-backed roadmap to survival and prosperity. By aggressively adopting salt-tolerant varieties, engineering the land to capture freshwater, diversifying crops, and embracing closed-loop integrated farming, coastal communities can turn their harshest vulnerabilities into strengths. The transition requires robust institutional backing, affordable technology, and active farmer participation. With these elements in place, the salty shores of West Bengal can be transformed from a landscape of climate vulnerability into a thriving model of agricultural resilience and enduring rural hope.

References

1. Ali, M. Y., & Hossain, M. E. (2019). Profiling climate smart agriculture for southern coastal region of Bangladesh and its impact on productivity, adaptation and mitigation. *EC Agriculture*, 5(9), 530-544.
2. Burman, D., Devika, S., Mandal, U. K., & Lama, T. D. (2026). Adoption of climate smart agricultural practices to mitigate multiple abiotic stresses in crops. In *Multiple Abiotic Stresses in Plants* (pp. 331-349). Academic Press.
3. Saha, Anubha, et al. "Qualitative evaluation of cropping system intensification in coastal saline zone of West Bengal, India." *J. Indian Soc. Coastal Agric. Res* 42.1 (2024): 147831.