



Biopesticides and Microbial Control: The Future of Sustainable Insect Pest Management

Pramod Kumar, Gaurav Kumar, Pushpraj, Shashibhushan Kumar, Bhanu Pratap,

*Raviprakash Kumar, Rohit Raj and Vikki Kumar Ram

M.Sc. Agriculture (Entomology), Department of Entomology, Late Chandrasekhar Ji

Purva Pradhan Mantri Smarak Mahavidyalaya, Rampur, Kanva, Saverai,

Ghazipur, Uttar Pradesh, India

*Corresponding Author's email: raviprakashkumarbth4@gmail.com

Insect pests remain a major constraint to crop productivity and farm profitability. Synthetic insecticides have provided effective and rapid pest suppression, but their repeated or inappropriate use can contribute to resistance, resurgence, residues and adverse effects on beneficial organisms and the environment. These concerns have increased interest in safer and more diversified pest-management approaches. Among these, **microbial biopesticides** have emerged as important tools for sustainable insect pest management. They use naturally occurring microorganisms or microbial agents that infect or otherwise suppress insect pests. The major groups include bacteria, entomopathogenic fungi, insect-pathogenic viruses and entomopathogenic nematodes. Microbial control does not aim to eliminate all insects from agricultural ecosystems. Instead, its objective is to suppress economically important pest populations while maintaining ecological balance. When combined with monitoring, economic thresholds, cultural practices, resistant varieties, natural enemies and selective insecticides, microbial biopesticides can become an important component of Integrated Pest Management (IPM).

Key message: Microbial biopesticides are most effective when they are used as part of a scientifically planned pest-management programme rather than as stand-alone substitutes for every conventional insecticide.

Major Microbial Groups Used in Pest Management

Table 1. Major microbial groups used in insect pest management

Microbial group	Important examples	Major target	Principal action
Bacteria	<i>Bacillus thuringiensis</i> (Bt)	Susceptible insect larvae	Gut disruption after ingestion
Entomopathogenic fungi	<i>Beauveria bassiana</i> , <i>Metarhizium anisopliae</i>	Various susceptible insects	Cuticle penetration and internal infection.
Viruses	NPV, GV	Mainly susceptible caterpillars	Infection and multiplication inside host cell
Nematodes	<i>Steinernema</i> , <i>Heterorhabditis</i>	Selected soil and cryptic pests	Host penetration and symbiotic bacteria

These groups differ considerably in their biology and application requirements. Therefore, the selection of a microbial agent should always be based on **pest species, developmental stage, crop, environmental conditions and application method**.

Bacterial Biopesticides: The Bt Success Story

Among microbial insecticides, *Bacillus thuringiensis* (Bt) is one of the most widely known and extensively used. Bt produces insecticidal proteins that are active against susceptible insects. When an appropriate Bt toxin is ingested by a susceptible larva, it acts primarily on the insect midgut, disrupting gut function and eventually causing mortality.




Figure 1. Simplified mode of action of *Bacillus thuringiensis* (Bt) against susceptible insect larvae.

Different Bt strains and toxins vary in their target ranges. Consequently, successful application depends on accurate identification of the target pest and selection of an appropriate Bt product. Bt illustrates an important principle of microbial pest management: specificity can be an advantage, but it also requires accurate pest identification.

Entomopathogenic Fungi: Infection Through the Insect Cuticle

Entomopathogenic fungi provide a different mechanism of biological control. *Beauveria bassiana* and *Metarhizium anisopliae* are important examples. Fungal spores attach to the insect body surface. Under suitable conditions, they germinate, penetrate the cuticle and develop inside the insect.




Figure 2. Infection process of entomopathogenic fungi through the insect cuticle.

Unlike microbial agents that must be ingested, these fungi can infect insects through the external body surface. This makes them useful against selected sucking and chewing pests. However, fungal performance is strongly influenced by **humidity, temperature, sunlight, rainfall and formulation quality**. Environmental conditions at the time of application can therefore determine whether a fungal product performs effectively in the field.

Insect-Pathogenic Viruses: Precision Through Specificity

Insect-pathogenic viruses, particularly **baculoviruses**, are valuable biological agents for controlling susceptible insect larvae. The important groups include **nucleopolyhedroviruses (NPVs)** and **granuloviruses (GVs)**. After entering a susceptible host, the virus multiplies within the insect and causes disease and mortality. Their high degree of host specificity can help reduce effects on many non-target organisms. However, this same specificity means that viral products cannot be expected to control unrelated pest species.

Successful viral control therefore depends on:

Correct pest identification + susceptible pest stage + appropriate timing + proper application

Baculoviruses are consequently most useful when the target pest and its biology are well understood.

Entomopathogenic Nematodes: Biological Control in the Soil

Some economically important insects spend part of their life cycle in soil or protected habitats, making them difficult to reach through conventional foliar applications.

Entomopathogenic nematodes (EPNs), particularly species belonging to *Steinernema* and *Heterorhabditis*, offer an important biological option for selected pests. Their infective juvenile stages locate insect hosts, enter them and release symbiotic bacteria. These bacteria contribute to infection and insect mortality.




Figure 3. Basic mechanism of entomopathogenic nematodes in insect hosts.

Their effectiveness depends strongly on soil moisture, temperature, soil characteristics and application conditions. Proper selection of nematode species and suitable timing is therefore essential.

Why Are Microbial Biopesticides Important?

Microbial agents can strengthen sustainable pest management in several ways.

Potential advantages

- They can provide alternatives to repeated use of conventional insecticides.
- Some microbial agents have relatively narrow host ranges.
- They can be incorporated into IPM programmes.
- Different biological modes of action can contribute to resistance-management strategies.
- Appropriate microbial agents can help conserve beneficial organisms and support ecological balance.
- They can be particularly useful where selective pest suppression is desirable.

However, **biological does not mean automatically effective**. Microbial products can fail when the wrong pest is targeted, when application timing is poor, or when environmental conditions are unsuitable. Therefore, the success of microbial control depends on matching **agent × pest × crop × environment × application method**.

Microbial Biopesticides and Chemical Insecticides

Microbial and chemical insecticides should not be considered mutually exclusive. Modern IPM uses different tools according to the pest situation.

Table 2. Key differences between microbial and conventional insecticides


Parameter	Microbial biopesticides	Conventional insecticides
Origin	Microorganisms or microbial products	Chemical active ingredients
Specificity	Often relatively specific	Variable
Speed of action	Often slower	Many provide rapid knockdown
Environmental sensitivity	Often high	Depends on product
Resistance	Possible	Important and widespread concern
IPM compatibility	Generally high	High when selectively used
Main requirement	Correct pest, timing and conditions	Correct pest, dose and timing


The choice should therefore be based on **pest biology and management objectives**, rather than simply on whether a product is biological or chemical.

From Laboratory Success to Field Performance

One of the greatest challenges in microbial pest management is maintaining efficacy under field conditions. Laboratory environments are relatively controlled, whereas agricultural fields experience changing temperature, humidity, sunlight, rainfall, wind, crop canopy and pest density. Microbial agents can therefore perform differently under different field conditions.

From discovery to field application





This process highlights why formulation and application technology are as important as discovering an effective microorganism.

Formulation and Application: The Critical Link

A microbial product must remain viable and effective during production, storage, transportation and application. It must also survive long enough in the field to reach and infect the target pest. Sunlight, temperature, humidity and rainfall can reduce the activity of some microbial agents. Formulation technologies therefore aim to improve **stability, shelf life, environmental protection and delivery to the target**.

Application timing is equally important.

A microbial product applied against the wrong pest stage or under unsuitable environmental conditions may provide poor control even when the microorganism itself is highly effective. This is why farmer training and technical guidance are essential for successful adoption.

Microbial Control Within IPM

Microbial biopesticides are most valuable when integrated with established principles of IPM.

A practical microbial-IPM sequence

- **Monitoring**

Regularly observe pest populations and crop condition.

- **Identification**

Confirm the pest species and developmental stage.

- **Assessment**

Determine whether the pest population has reached an economically important level.

- **Selection**

Choose the most appropriate management option.

- **Application**

Apply the microbial product at the correct time and according to the recommended method.

- **Evaluation**

Assess pest suppression and crop response.

- **Continued monitoring**

Determine whether additional intervention is required.

Finding an insect in a crop does not automatically mean that pesticide application is necessary.

Economic thresholds, natural enemies, crop stage and pest biology should all be considered before intervention.

Opportunities for India

India has considerable potential for microbial pest management because of its diverse crops, agro-climatic conditions and wide range of insect pests.

Microbial agents such as *Bacillus thuringiensis*, *Beauveria bassiana*, *Metarhizium anisopliae* and baculoviruses have received substantial research attention.

However, wider adoption requires more than product availability. Important requirements include:

- reliable product quality,
- appropriate storage,
- correct pest identification,
- proper application timing,
- farmer training,

- field validation,
- quality assurance, and
- compliance with applicable regulations.

For small and marginal farmers, affordability and accessibility are especially important.

Agricultural universities, research institutions, extension agencies, Krishi Vigyan Kendras and trained service providers can play an important role in transferring microbial technologies from research programmes to farm-level practice.

Microbial Control Meets Precision Agriculture

The next stage of microbial pest management may involve integration with digital agriculture. Smart insect traps, weather stations, drones, remote sensing and AI-based image analysis can improve pest monitoring and help identify areas requiring attention.



Figure 4. Precision-agriculture workflow for targeted microbial pest management.

Such systems could help shift pest management from routine calendar-based spraying toward **more targeted and information-based intervention**.

However, digital tools should support rather than replace biological expertise.

AI can identify patterns; entomology provides biological interpretation.

Climate Change and Microbial Control

Changing climatic conditions may influence both insect populations and microbial agents.

Temperature, humidity and rainfall affect insect development as well as the survival and infectivity of microbial pathogens. Changes in pest distribution and crop-growing seasons may also alter when and where microbial products can be used effectively.

Future research should therefore emphasize:

- environmentally robust microbial strains,
- improved formulations,
- better environmental forecasting,
- optimized application methods, and
- region-specific field validation.

Developing microbial technologies that remain effective under variable field conditions will be essential for their wider adoption.

Major Challenges

Despite their promise, microbial biopesticides face several limitations:

Variable field efficacy

Environmental conditions can substantially influence performance.

Formulation and shelf life

Microbial agents require appropriate stabilization and storage.

Application knowledge

Incorrect pest identification, timing or application can reduce efficacy.

Production and quality control

Commercial products require consistent microbial viability and quality.

Resistance management

Microbial agents can also face reduced sensitivity, making diversified management important.

Regulatory requirements

Products must meet applicable safety, efficacy and registration standards.

These challenges require collaboration among **entomologists, microbiologists, agronomists, industry, extension personnel and farmers.**

The Future: Toward Smarter Biological Pest Management

The future of microbial control is unlikely to depend on a single “miracle” organism. Greater progress is likely to come from combining biological control with modern monitoring and decision-support technologies.

A future pest-management system could integrate:



Figure 5. Integrated framework for smarter biological pest management

Such integration could make pest management more timely, targeted and environmentally responsible. The key objective should not be to maximize pesticide use with better technology. It should be to **improve the quality of pest-management decisions.**

Conclusion

Microbial biopesticides represent an important opportunity for sustainable insect pest management. Bacteria, fungi, viruses and entomopathogenic nematodes provide different biological mechanisms for suppressing economically important pests. Their greatest value lies not in completely replacing conventional insecticides, but in **diversifying the pest-management toolbox.** Successful microbial control requires accurate pest identification, suitable microbial agents, quality formulations, appropriate timing, favourable environmental conditions and integration with IPM. The future of sustainable crop protection will therefore depend on combining **biology, entomology, technology and farmer knowledge.** Microbial control offers a practical step toward this future—where insect pests are monitored carefully, interventions are targeted, beneficial organisms are conserved and crop protection becomes increasingly precise. **The goal is not to eliminate every insect. The goal is to manage pests intelligently, sustainably and only when necessary.**

References

1. Lacey, L. A., Grzywacz, D., Shapiro-Ilan, D. I., Frutos, R., Brownbridge, M., & Goettel, M. S. (2015). Insect pathogens as biological control agents: Back to the future. *Journal of Invertebrate Pathology*, 132, 1–41. <https://doi.org/10.1016/j.jip.2015.07.009>
2. Kumar, K. K., Sridhar, J., Murali-Baskaran, R. K., Senthil-Nathan, S., Kaushal, P., Dara, S. K., & Arthurs, S. (2019). Microbial biopesticides for insect pest management in India: Current status and future prospects. *Journal of Invertebrate Pathology*, 165, 74–81. <https://doi.org/10.1016/j.jip.2018.10.008>
3. Chandler, D., Bailey, A. S., Tatchell, G. M., Davidson, G., Greaves, J., & Grant, W. P. (2011). The development, regulation and use of biopesticides for integrated pest management. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366(1573), 1987–1998. <https://doi.org/10.1098/rstb.2010.0390>
4. Glare, T., Caradus, J., Gelernter, W., Jackson, T., Keyhani, N., Köhl, J., Marrone, P., Morin, L., & Stewart, A. (2012). Have biopesticides come of age? *Trends in Biotechnology*, 30(5), 250–258. <https://doi.org/10.1016/j.tibtech.2012.01.001>
5. Faria, M. R. de, & Wraight, S. P. (2007). Mycoinsecticides and mycoacaricides: A comprehensive list with worldwide coverage and international classification of formulation types. *Biological Control*, 43(3), 237–256. <https://doi.org/10.1016/j.biocontrol.2007.08.001>
6. van Lenteren, J. C., Bolckmans, K., Köhl, J., Ravensberg, W. J., & Urbaneja, A. (2018). Biological control using invertebrates and microorganisms: Plenty of new opportunities. *BioControl*, 63, 39–59. <https://doi.org/10.1007/s10526-017-9801-4>
7. Food and Agriculture Organization of the United Nations (FAO). Integrated Pest Management (IPM). FAO Plant Production and Protection Division.
8. Food and Agriculture Organization of the United Nations (FAO). Introduction: Biological Pest Control Agents. Pesticide Registration Toolkit