



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 10 (October, 2026)

Available online at <http://www.agrimagazine.in>

©Agri Magazine, ISSN: 3048-8656

Regenerative Vegetable Farming: Practical Approaches for Soil Health and Resilient Production

*Choudhary Snehal

Postgraduate Scholar, Department of Vegetable Science, Mahatma Phule Krushi Vidyapeeth (MPKV), Rahuri, Maharashtra-413722, India

*Corresponding Author's email: snehalchowdhary020@gmail.com

Regenerative vegetable farming applies soil-centred practices to restore soil health, support biodiversity and sustain vegetable production. Its central principles are to reduce unnecessary soil disturbance, keep soil covered, maintain living roots for as much of the year as possible and diversify crops. Practical options include cover crops, crop rotation, compost and farmyard manure, residue mulching, reduced tillage, deep-rooted crops, biochar where locally suitable, and integrated pest management. These practices can improve soil structure, water infiltration, moisture retention and biological activity while reducing erosion and dependence on purchased inputs. Outcomes depend on soil type, climate, crop choice, management and the length of time practices are maintained. Regenerative methods can be introduced gradually on vegetable farms by starting with soil cover, locally available organic materials and planned rotations, then monitoring crop performance, labour, input costs and soil condition. This approach offers a practical pathway towards more resilient vegetable production, although local trials and economic assessment are needed to identify the most effective combinations for each farming system.

Keywords: regenerative agriculture; vegetable farming; soil organic carbon; cover crops; mulching; crop rotation; water management

Introduction

Vegetable production is increasingly challenged by soil degradation, erosion, irregular rainfall, rising input costs and climate variability. Repeated intensive tillage, limited crop diversity and the removal or burning of crop residues can weaken soil structure and reduce the organic matter available to soil organisms. These problems may affect water infiltration, moisture storage, nutrient cycling and the long-term productivity of vegetable fields. Regenerative vegetable farming adapts the principles of regenerative agriculture to vegetable-based systems. Rather than focusing only on maintaining current production, it seeks to improve the condition and functioning of the farm ecosystem over time. Soil cover, living roots, crop diversity, organic amendments and reduced disturbance are used to support soil organisms and protect the soil surface. Permanent or semi-permanent beds, where appropriate, can help organise repeated vegetable crops while limiting unnecessary disturbance between plantings. Regenerative agriculture can also contribute to diversified diets and farm livelihoods when it supports reliable production of a range of crops. However, it should not be treated as a single fixed package. Practices need to be selected according to local climate, soil, irrigation facilities, labour availability, market demand and the resources accessible to farmers.

Principles of Regenerative Vegetable Farming

The main principles are:

1. Minimise soil disturbance: Reduce unnecessary tillage and avoid working soil when it is excessively wet. Reduced disturbance can help preserve soil aggregates and biological habitats.
2. Keep the soil covered: Use crop residues, suitable organic mulches or cover crops to protect the soil from raindrop impact, erosion and direct sunlight.
3. Maintain living roots: Plan rotations and cover crops so that living roots remain in the soil for as much of the year as practical, without competing excessively with the vegetable crop for water.
4. Increase crop diversity: Rotate crops from different botanical families and include legumes or other suitable cover crops to diversify rooting patterns and residue inputs.
5. Recycle farm resources: Use safe, mature compost and well-decomposed farmyard manure according to crop and soil needs. Where livestock are integrated, manage grazing and manure to avoid compaction, contamination and nutrient losses.

Practical Methods for Vegetable Farms

Cover crops and green manures: Cover crops can be grown between vegetable crops or during periods when beds would otherwise remain bare. Suitable legumes, grasses and cruciferous species can provide ground cover, add biomass, suppress some weeds and capture nutrients that might otherwise be lost. Legumes may contribute nitrogen through biological fixation, but species selection and termination timing are important. Cover crops should be managed so they do not deplete water needed by the next vegetable crop or become difficult to control.

Crop rotation and diversification: Avoid planting the same vegetable family repeatedly in the same bed. Rotating crops with different nutrient requirements and rooting patterns can help disrupt some pest and disease cycles and diversify residue inputs. For example, a farm may alternate solanaceous vegetables such as tomato with crops from other families, selected according to local suitability and market needs. Rotation plans should also consider the host range of important soil-borne pests and pathogens.

Mulching and residue retention: Apply clean crop residues or suitable organic mulch to reduce soil exposure, moderate soil temperature, conserve moisture and suppress weeds. Keep mulch away from plant stems where excessive moisture could encourage disease. Use only materials that are safe for food production and free from persistent herbicide residues or contaminants. In areas where residues are needed for livestock feed or fuel, farmers may need to balance these competing uses.

Compost and farmyard manure: Mature compost and well-decomposed manure can add organic matter and nutrients. Application rates should be guided by soil testing, crop nutrient demand and the nutrient content of the material. Excessive application can cause nutrient imbalances or losses to water. Fresh manure should not be applied in ways that create food-safety risks, particularly close to harvest.

Reduced tillage and permanent beds: Where soil and equipment permit, reduce the frequency and depth of tillage. Permanent beds can concentrate traffic in paths and reduce compaction in the growing area. Small-scale farmers may begin by reducing repeated bed preparation rather than immediately adopting a strict no-till system. Transplanting equipment, weed management and the condition of the soil must be considered before changing the tillage system.

Deep-rooted crops and biological activity: Suitable deep-rooted cover crops can explore lower soil layers, take up nutrients and create channels that may help subsequent roots. Diverse root systems and organic inputs provide food and habitat for soil organisms. Benefits depend on species, soil depth, moisture and management; therefore, crop mixes should be tested locally.

Biochar: Biochar is a carbon-rich material produced by heating biomass under limited oxygen. Depending on its source and production quality, it may improve some soil properties and retain nutrients or water. Its performance varies with soil type, feedstock and application

rate. Use only well-characterised, uncontaminated biochar and evaluate cost-effectiveness before large-scale application.

Integrated pest and weed management: Combine crop rotation, field sanitation, mulching, monitoring, traps, conservation of natural enemies and locally approved biopesticides when needed. Neem-based products or plant-derived preparations should be used only where appropriate and according to product guidance and local regulations. Regenerative management does not mean ignoring serious pest outbreaks; timely, evidence-based control is still essential.

Benefits And Implementation Challenges

When sustained and appropriately managed, regenerative practices can improve soil aggregation, infiltration, moisture retention and biological activity. Cover crops and residue retention protect the soil surface, while organic amendments supply carbon and nutrients. These changes may reduce erosion and help crops cope with short dry periods. Increasing crop diversity can also support pollinators and other beneficial organisms when flowering resources and suitable habitats are available. Soil organic carbon is an important indicator of soil condition and is relevant to climate-change mitigation. However, carbon accumulation is not guaranteed, and rates vary with climate, soil texture, previous land use, inputs and management duration. Claims about carbon sequestration should therefore be interpreted in context and, where possible, supported by repeated soil measurements using consistent sampling methods. The transition can involve additional planning, labour for mulch or compost handling, cover-crop seed costs and short-term uncertainty in yields. Cover crops may compete for water, organic materials may be scarce, and reduced tillage can initially complicate weed control. Organic inputs also need careful nutrient management. For these reasons, farmers should avoid changing every practice at once.

A practical starting plan is to select one or two beds or a small field section; record the current crop, yield, labour and input costs; introduce soil cover and a suitable rotation; and compare the results with a conventionally managed section. Monitor crop establishment, weeds, pest incidence, irrigation use, yield, marketable produce and total costs over more than one season. Soil organic matter or soil organic carbon, infiltration and bulk density can be assessed periodically where facilities are available. Expand the practices that demonstrate agronomic and economic value under local conditions. Livestock integration and agroforestry may be useful where farm size, water, fencing, labour and markets permit. Manure must be managed safely, while trees should be selected and placed to minimise unwanted shade and competition with vegetable crops. Government support, farmer training, local demonstrations and market access can help reduce transition risks.

Conclusion

Regenerative vegetable farming offers a practical framework for improving soil condition and building resilience through reduced soil disturbance, continuous soil cover, living roots, crop diversification and careful recycling of organic resources. Cover crops, rotations, mulching, compost, reduced tillage and integrated pest management can be introduced gradually and adapted to local vegetable systems. Their benefits depend on sound crop planning, appropriate nutrient and water management, and consistent monitoring. Because costs, yields and soil responses vary among farms, farmers should test practices on a manageable scale and assess both environmental and economic outcomes before expanding them. Long-term, location-specific trials are needed to identify the most effective combinations for different Indian agro-climatic regions.

References

1. Anderson, M. D. and Rivera-Ferre, M. (2021). Food system narratives to end hunger: Extractive versus regenerative. *Current Opinion in Environmental Sustainability*, 49, 18–25.

2. Avasiloaiei, D. I., Calara, M., Brezeanu, M. P. and Cristea, T. O. (2023). Review on the sustainability of some regenerative agriculture practices for organic vegetable growing. *Scientific Papers Series B, Horticulture*, 67(2), 189–198.
3. Diwan, A. D., Harke, S. N., Pande, B. N. and Panche, A. (2021). Regenerative agriculture farming. *Indian Farming*, 71(12), 42–45.
4. Gelaw, A. M., Singh, B. R. and Lal, R. (2014). Soil organic carbon and total nitrogen stocks under different land uses in a semi-arid watershed in Tigray, Northern Ethiopia. *Agriculture, Ecosystems & Environment*, 188, 256–263.
5. Lal, R. (2004). Soil carbon sequestration impacts on global climate change and food security. *Science*, 304(5677), 1623–1627.
6. Liao, Y., Wu, W. L., Meng, F. Q., Smith, P. and Lal, R. (2015). Increase in soil organic carbon by agricultural intensification in northern China. *Biogeosciences*, 12(5), 1403–1413.
7. Marichamy, M. S. and Kanthaswamy, V. (2024). Natural farming: Embracing regenerative agriculture for sustainable crop production. *Journal of Experimental Agriculture International*, 46(8), 855–865.
8. Matsuura, E., Komatsuzaki, M. and Hashimi, R. (2018). Assessment of soil organic carbon storage in vegetable farms using different farming practices in the Kanto Region of Japan. *Sustainability*, 10(1).
9. Patil, M., Perumal, C. and Choudhari, P. (2025). Differential impacts of regenerative agriculture practices on soil organic carbon: a meta-analysis of studies from India. *Scientific Reports*, 15, 33470.
10. Rempelos, L., Kabourakis, E. and Leifert, C. (2023). Innovative organic and regenerative agricultural production. *Agronomy*, 13(5).
11. Rice, S. W. (2025). Land transitions to regenerative vegetable farming: Assessing opportunities for established traditional farms and small vegetable growers [Doctoral dissertation, Harvard University].