



## Beneath Our Feet: Unravelling the Hidden Burden of Heavy Metal Contamination

\*Kancha Sai Kiran and P. Elavarasi

Department of Soil Science and Agricultural Chemistry, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore-641003, India

\*Corresponding Author's email: [saikiran.phdsac2023@tnau.ac.in](mailto:saikiran.phdsac2023@tnau.ac.in)

A green field often gives us a reassuring picture of healthy soil. Crops grow, leaves appear green, and harvests fill our markets. Yet, beneath this apparently healthy surface, another story may be unfolding. Soil can silently accumulate contaminants that are invisible to the eye but capable of affecting soil functions, crop productivity, food quality and environmental health. Among the contaminants of growing concern are potentially toxic elements such as cadmium (Cd), lead (Pb), arsenic (As), chromium (Cr), nickel (Ni), mercury (Hg), copper (Cu) and zinc (Zn). Some of these elements, particularly Cu and Zn, are essential micronutrients required by plants in small quantities. However, when their concentrations become excessive, they can become toxic. Others, such as Cd and Pb, have no known essential nutritional role in plants and can pose risks even when present in relatively low concentrations. The challenge is not simply that these elements are present in soil. The real concern is their persistence, mobility, bioavailability and movement through the soil-plant system. Unlike many organic pollutants, metals and metalloids cannot be destroyed through ordinary biological degradation. They can change their chemical forms, move between different soil pools and remain in the environment for long periods. This makes heavy metal contamination a particularly important issue for agricultural soils, where the quality of soil is directly connected to the quality and safety of food.

### Where do these hidden contaminants come from?

Heavy metals in soil may originate from both natural geological processes and human activities. The weathering of parent rocks naturally contributes metals to soils. However, human activities can significantly increase their concentrations and alter their distribution. Mining and mineral processing, metal smelting, industrial emissions, traffic-related deposition, sewage and wastewater irrigation, improper waste disposal and certain agricultural inputs are important sources of metal contamination. Contaminated irrigation water can introduce metals repeatedly into agricultural land, while atmospheric deposition can transport contaminants from industrial and urban areas to surrounding soils. The problem may become more serious when contamination occurs repeatedly over many years. A single input may not produce an immediately visible effect, but continuous additions can gradually increase the soil burden. The FAO and UNEP have highlighted soil pollution as a major threat to soil health because contaminants can compromise food production, water quality, biodiversity and ecosystem functions. Importantly, soil pollution is often invisible, meaning that contamination may remain unnoticed until its effects become significant. Thus, the question is not simply *"How did the metal enter the soil?"* but also *"What happens to it after it enters?"*


### The soil is not a storage box

It is tempting to imagine soil as a container that simply stores whatever contaminants enter it. In reality, soil is a highly dynamic chemical and biological system. Once metals enter soil,

they can become associated with clay minerals, soil organic matter, iron and manganese oxides, carbonates and other soil constituents. They may also remain dissolved in the soil solution or occur in different chemical forms. These forms determine how easily a metal can move through soil, be taken up by plants or reach groundwater. This is where the concept of bioavailability becomes extremely important. A soil may contain a high total concentration of a particular metal, but if most of that metal is strongly bound to soil components, its immediate availability to plants may be comparatively low. Conversely, a smaller total concentration may still present a concern if a large fraction occurs in a readily available form. Several soil properties influence this behaviour, including soil pH, organic matter, clay content, redox conditions, mineral composition, moisture conditions and competing ions. Consequently, two soils containing similar total concentrations of a metal may not necessarily present the same level of risk to crops. This distinction is fundamental to modern soil pollution assessment. The important question is not only “How much metal is in the soil?” but also “How much of it is available to move, leach or enter plants?”

### When contamination meets the root zone

The rhizosphere-the narrow region of soil influenced by plant roots-is a hotspot of chemical and biological activity. Roots release organic compounds, interact with microorganisms and continuously exchange nutrients and ions with the surrounding soil solution. Under favourable conditions, potentially toxic elements can become available for root uptake. Once absorbed, their movement within the plant depends on the element, plant species, root characteristics, environmental conditions and physiological mechanisms. For example, excessive concentrations of Cd, Pb or other toxic elements can interfere with nutrient uptake, root development, photosynthesis and cellular metabolism. Metal stress can also promote the formation of reactive oxygen species (ROS), resulting in oxidative stress and damage to cellular components. However, plants do not respond identically to all metals. Some species can tolerate relatively high concentrations by restricting uptake, binding metals within roots or storing them in less metabolically active cellular compartments. Other species may absorb and translocate particular elements more readily. Therefore, soil contamination does not automatically translate into identical contamination of every crop. Crop species, cultivar, growth stage, soil conditions and the chemical form of the contaminant all influence the soil-to-plant transfer process.



## From soil to food: when the hidden burden becomes a public concern


Agricultural soil occupies a special position in the contamination pathway because it can connect environmental pollution with the food chain.

A simplified pathway can be represented as:

**Contaminated source → Soil → Soil solution → Plant roots → Plant tissues → Food → Consumer**

When metals become sufficiently available for plant uptake, they may accumulate in roots, stems, leaves, fruits, grains or other edible tissues. The extent of accumulation varies considerably among elements and crops. Rice, vegetables and other food crops grown in contaminated environments have received particular scientific attention because certain metal(loid)s can enter edible tissues. Research on contaminated agricultural systems has demonstrated that understanding metal mobility, speciation and bioavailability is essential for evaluating potential human-health risks. Importantly, the total concentration of a metal in soil alone cannot reliably predict its concentration in a particular crop. The relationship between soil metal concentration and plant accumulation is influenced by numerous soil and plant factors and may not be linear. This is why soil testing, crop monitoring and risk assessment need to work together.

**The unseen impact on soil life** Heavy metal contamination is not only a problem for plants. Soil itself is a living ecosystem. Bacteria, fungi, actinomycetes, earthworms and other soil organisms contribute to decomposition, nutrient cycling, aggregate formation and maintenance of soil fertility. Elevated concentrations of potentially toxic elements can disturb microbial communities and biological processes, depending on the element, concentration and soil conditions. Changes in microbial activity can have consequences beyond microorganisms themselves. If important biological processes involved in carbon and nutrient cycling are disrupted, the functional capacity of soil may gradually decline. Therefore, heavy metal contamination should not be viewed simply as a chemical problem. It is a chemical, biological and ecological problem simultaneously. Recent research has emphasized the links between heavy metal contamination, soil properties, microorganisms, plant growth and food-chain transfer. Can we clean contaminated soil? The answer is yes-but there is no single solution suitable for every contaminated soil. The appropriate strategy depends on the type and concentration of contaminant, soil properties, land use, contamination depth, crop requirements and the intended end use of the land.



**1. Immobilisation: making metals less available**

Immobilisation aims to reduce the mobility and bioavailability of metals rather than physically removing them from the soil. Materials such as clay minerals, lime-containing amendments, biochar and other sorbent materials can interact with metals and reduce their availability under suitable conditions. Changes in soil pH, adsorption, precipitation and complexation are among the mechanisms involved. For example, clay minerals such as bentonite, sepiolite and palygorskite have been investigated for reducing the bioavailability of metals in contaminated agricultural soils. However, immobilisation does not mean that the contaminant has disappeared. The metal remains in the soil, and its long-term stability must be monitored.

**2. Phytoremediation: allowing plants to assist**

Plants can also become part of the solution. Phytoremediation uses plants to remove, stabilize or otherwise manage contaminants in soil. Some plant species can tolerate and accumulate particular metals, while others can help stabilize contaminants and reduce their movement. This approach can be environmentally attractive, but its effectiveness depends strongly on plant species, contaminant type, soil conditions and the duration of treatment. Plant biomass containing accumulated metals must also be managed carefully after harvest.

**3. Microbial and biological approaches**

Microorganisms can influence metal behaviour through processes such as biosorption, bioprecipitation, transformation and changes in the soil chemical environment. Rhizosphere microorganisms may also interact with plants and influence metal uptake. Biological remediation is therefore receiving increasing attention as a potentially sustainable component of integrated soil restoration. However, field conditions are considerably more complex than controlled laboratory systems, and long-term effectiveness must be demonstrated before widespread application.

**4. Integrated soil management**

Increasingly, researchers are moving away from the idea that one technology can solve every contamination problem. A combination of soil amendments, appropriate crop selection, irrigation management, nutrient management, phytoremediation and biological approaches may provide better results in certain situations. Recent research emphasizes that remediation should be selected according to contaminant characteristics, soil conditions and risk pathways rather than applying the same treatment universally.

**Prevention is better than remediation**

Remediating contaminated soil can be technically difficult, expensive and time-consuming. Therefore, preventing contamination at its source is often more effective than attempting to restore severely contaminated soil later.

Agricultural soils should be protected through:

- Regular monitoring of soil and irrigation-water quality.
- Responsible use of fertilizers, pesticides and organic amendments.
- Proper management of industrial and municipal wastes.
- Avoidance of untreated or poorly characterized wastewater for irrigation.
- Testing manure, compost and other amendments where contamination is a concern.
- Monitoring potentially contaminated agricultural areas near industries, mining operations and waste-disposal sites.
- Periodic assessment of metal concentrations and their bioavailable fractions where contamination is suspected.

Particular attention should be given to repeated applications of organic amendments. Organic materials can improve soil fertility and soil structure, but their quality and contaminant content should be assessed before long-term application. Recent research also emphasizes that manure can contain metals such as Cu, Zn, Cd, Pb and Cr depending on its source and management history. Good soil management therefore means not only feeding the soil, but also protecting what enters the soil.

## Looking beneath the surface

Heavy metal contamination reminds us that soil health cannot be judged by appearance alone. A field may be green. A crop may look vigorous. The soil may appear dark and fertile. Yet chemical contaminants may be accumulating quietly beneath the surface. This is why modern soil science must look beyond total metal concentrations and consider the complete **soil-metal-microbe-plant continuum**. Understanding the chemical forms of metals, their interaction with soil minerals and organic matter, their mobility, their bioavailability and their transfer into crops provides a much clearer picture of environmental risk. The goal should not simply be to remove every trace of a naturally occurring element. Instead, the focus should be on preventing harmful accumulation, minimizing bioavailable and mobile fractions, protecting soil functions, preventing food-chain transfer and ensuring long-term soil safety.

## The future lies beneath our feet

The soil beneath our feet is much more than a medium in which plants grow. It is a living interface connecting geology, water, microorganisms, plants, agriculture and human society. Heavy metal contamination challenges us because it is persistent and often invisible. But it also demonstrates why soil science is so important. Through careful monitoring, risk assessment, improved agricultural practices and scientifically validated remediation technologies, contaminated soils can be managed more responsibly.

## References

1. FAO & UNEP. (2021). *Global Assessment of Soil Pollution: Report*. Food and Agriculture Organization of the United Nations and United Nations Environment Programme.
2. Mai, X., Tang, J., Tang, J., Zhu, X., Yang, Z., Liu, X., Zhuang, X., Feng, G., & Tang, L. (2025). Research progress on the environmental risk assessment and remediation technologies of heavy metal pollution in agricultural soil. *Journal of Environmental Sciences*, 149, 1–20.