AGRI MAGAZINE
(International E-Magazine for Agricultural Articles)
Volume: 03, Issue: 10 (October, 2026)
Available online at http://www.agrimagazine.in
©Agri Magazine, ISSN: 3048-8656

Ponnagani Keerai (*Alternanthera sessilis*): Nutritional Importance, Traditional Value, Cultivation Potential and Future Prospects

*Gokul MK, Karthick C, Mussamil and Gokulavani
Department of Horticulture, Faculty of Agriculture, Annamalai University, Tamil Nadu, India
*Corresponding Author's email: gokiiikannan1009@gmail.com

Ponnagani keerai, commonly known as sessile joyweed and botanically identified as Alternanthera sessilis, is an edible leafy vegetable widely used in traditional food systems. It is valued particularly as a locally available leafy green and is suitable for repeated harvesting. Like other leafy vegetables, it can contribute vitamins, minerals, dietary fibre and bioactive plant compounds to the diet. Its ability to grow under warm and humid conditions and its simple propagation make it a useful crop for kitchen gardens and small-scale vegetable production.



Figure 1. Ponnagani keerai (*Alternanthera sessilis*)

Botanical Description and Plant Characteristics

Alternanthera sessilis is a herbaceous plant with soft, branched stems that may root at the nodes when they come in contact with moist soil. The leaves are simple, opposite, tender and generally green. Small flowers occur in compact clusters in the leaf axils. The plant produces tender shoots and leaves that are commonly harvested for culinary use. Its creeping or spreading growth habit allows rapid establishment under suitable moisture conditions.

Nutritional and Functional Importance

Ponnagani keerai can form part of a diversified diet based on leafy vegetables. Leafy greens are important sources of dietary fibre and micronutrients, while their green tissues contain pigments and other phytochemicals. The nutritional composition of Ponnagani can vary with cultivar or accession, maturity, growing environment and method of preparation. Traditional claims about medicinal benefits should be distinguished from scientifically established health effects; further nutritional and clinical research can help characterize its functional potential.

Traditional and Culinary Importance

Ponnagani keerai has a long association with traditional food practices in parts of South India. Tender leaves and shoots are generally cooked as a leafy vegetable and may be incorporated into poriyal, kootu, curry, soups or other preparations. Its acceptance in traditional cuisine provides an opportunity to promote the crop through recipe development, consumer education and local food enterprises.

Cultivation Potential

The crop can be established using tender stem cuttings or other suitable planting material. A well-drained fertile soil with adequate organic matter and regular moisture supports vigorous vegetative growth. Land preparation should provide a fine tilth, followed by planting of

healthy cuttings at a convenient spacing. Organic manures and balanced nutrient management can support leaf production. Regular irrigation is important, particularly during dry periods. Weed management during early establishment and monitoring for pests and diseases help maintain quality produce.

Harvesting and Yield Potential

Tender shoots and leaves can be harvested after adequate vegetative establishment. Repeated harvesting may encourage production of new tender shoots when adequate moisture and nutrients are maintained. Harvesting should preferably be done during cooler parts of the day and the produce should be handled carefully to reduce wilting and mechanical damage.

Socio-economic Importance

Ponnagani can be useful for household nutrition as well as small-scale commercial vegetable production. Its short harvesting cycle and potential for repeated picking can provide opportunities for frequent local sales. Because it is a traditional leafy vegetable, strengthening local markets, improving hygienic packing and developing value-added products may increase its commercial visibility.

Post-harvest Handling and Value Addition

Fresh Ponnagani leaves are highly perishable and therefore benefit from rapid marketing, shade handling and appropriate cooling. Washing should use clean water and excessive mechanical handling should be avoided. Potential value-added products include cleaned and packed fresh greens, ready-to-cook leafy vegetable packs, dried leafy products and incorporation into traditional food preparations. Processing methods should be evaluated for their effects on nutrients, colour, flavour and shelf life.

Constraints

Despite its traditional importance, Ponnagani is less commercially organized than major leafy vegetables. Important constraints include limited availability of standardized planting material, inadequate crop-specific production recommendations, short post-harvest life, inconsistent market demand and limited organized value addition. More research is also needed on varietal characterization, nutrition, pest and disease management, production economics and post-harvest technology.

Future Prospects

Greater attention to indigenous leafy vegetables can support dietary diversification and conservation of crop diversity. Future work on Ponnagani should include collection and characterization of diverse accessions, selection for tender leaf quality and yield, standardized agronomic practices, nutrient profiling, safe production systems, improved packaging and development of convenient food products. Farmer groups, local markets, extension agencies and small food enterprises can help connect production with consumers.

Conclusion

Ponnagani keerai is a traditional leafy vegetable with nutritional, culinary and livelihood value. Its simple growth habit, repeated harvesting potential and familiarity in traditional food systems provide a foundation for wider utilization. Scientific research, improved planting material, better post-harvest handling, consumer awareness and value addition can contribute to its conservation and future commercialization.

Table 1. Major areas for development of Ponnagani keerai

Area	Current importance	Future opportunity
Nutrition	Traditional leafy green	Detailed nutrient and phytochemical profiling
Production	Suitable for small-scale cultivation	Standardized agronomic practices and planting material
Harvesting	Tender leaves and shoots	Improved repeated-harvest management

Area	Current importance	Future opportunity
Marketing	Mostly local consumption	Organized local markets and hygienic packing
Value addition	Mainly fresh culinary use	Ready-to-cook and dried products

References

1. Ali, A., & Bhattacharjee, B. (2023). Nutrition security, constraints, and agro-diversification strategies of neglected and underutilized crops to fight global hidden hunger. *Frontiers in Nutrition*.
2. Ebert, A. W. (2014). Potential of underutilized traditional vegetables and legume crops to contribute to food and nutritional security, income and more sustainable production systems. *Sustainability*, 6(1), 319–335.
3. Mayes, S., Massawe, F. J., Alderson, P. G., Roberts, J. A., Azam-Ali, S. N., & Hermann, M. (2012). Food security: the potential for underutilized crops to improve security of food production. *Journal of Experimental Botany*, 63(3), 1075–1079.
4. Savita, Rawat, M., & Vimal, V. (2023). Underutilized vegetables — introduction and identification. In Underutilized Vegetable Crops. Springer.