



AGRI MAGAZINE

(International E-Magazine for Agricultural Articles)

Volume: 03, Issue: 10 (October, 2026)

Available online at <http://www.agrimagazine.in>

©Agri Magazine, ISSN: 3048-8656

Major Reproductive Problems in Goat Farming

*Mote Sujit

Livestock Production and Management Section, Indian Veterinary Research Institute,
Izatnagar, Uttar Pradesh, India

*Corresponding Author's email: sujitmote0@gmail.com

Goat farming plays a significant role in rural livelihoods by providing meat, milk, skin, manure, and income, particularly for small and marginal farmers. However, reproductive disorders and pregnancy-related complications can adversely affect productivity, reproductive efficiency, and farm profitability. Major reproductive problems in goats include dystocia, uterine prolapse, retention of fetal membranes, hydrometra or pseudopregnancy, ovarian cysts, and abortion. Dystocia may arise from maternal factors, inadequate cervical dilation, or abnormal fetal presentation. Nutritional deficiencies, particularly of iodine, manganese, vitamin A, and energy, may impair reproductive performance and increase pregnancy losses. Infectious agents, including *Chlamydia abortus*, *Listeria monocytogenes*, *Salmonella* spp., and *Toxoplasma gondii*, are also associated with abortion and reproductive failure. Early diagnosis, appropriate veterinary treatment, balanced nutrition, hygienic kidding management, and effective disease prevention are essential for reducing reproductive losses. Improved reproductive health management can enhance kidding rates, kid survival, animal welfare, and the overall profitability of goat farming enterprises.

Keywords: Goat farming, Reproductive disorders, Dystocia, Hydrometra, Abortion, Uterine prolapse, Reproductive health.

Introduction

Goat farming offers significant opportunities for rural development because it provides rural farmers with meat, milk, skin, manure, and wool, thereby strengthening their economic security. Goat products also have export potential, contribute to capital accumulation, and provide household income, employment, and nutritional security. Compared with other livestock enterprises, goat farming offers greater economic and managerial advantages because it requires low initial investment, fewer inputs, has a shorter generation interval, and allows relatively easy marketing. Goat farming can provide year-round self-employment to small and marginal farmers without significantly affecting their primary occupation. In the absence of major infectious causes, infertility is generally not a major problem in goats.

Dystocia (Difficult Parturition)

The first stage of labour lasts approximately 6–12 hours, while the second stage lasts 0.5–1 hour. Vocalisation may be observed in goats during parturition. The incidence of dystocia varies considerably, ranging from approximately 8% to 50% in different reports. Failure of the cervix to dilate is a major cause of dystocia and may subsequently lead to uterine inertia. Abnormal lateral deviation of the head and flexion of the carpal and hock joints are common fetal causes of dystocia. These abnormalities may increase the apparent size of the fetus and obstruct delivery. Dystocia can be managed through correction of fetal malpresentation or by caesarean section. However, in sheep and goats, partial extraction of the fetus through the cervix may sometimes be possible only after adequate cervical dilation. The birth canal of a doe in labour is delicate, and excessive force used to extract a malformed or oversized fetus

can result in uterine rupture, particularly when fetal abdominal organs protrude. Therefore, careful handling is essential during manual delivery. In some regions of Uttar Pradesh, dystocia associated with fetal goitre has been reported. Iodine deficiency is considered a possible underlying cause. To prevent this condition, pregnant animals should receive a properly balanced ration containing all essential nutrients. Iodine-deficient pregnant goats should receive an appropriate iodine supplement, such as iodised salt. The text specifies an iodine concentration of 0.019% in salt.

Prolapse

Prolapse can be defined as the displacement of an organ or part of an organ from its normal anatomical position. In goats, uterine prolapse occurs when the cervix is open and the uterus protrudes through the vagina. It generally appears as a large mass of tissue emerging from the birth canal and usually occurs during the third stage of labour, after the fetus has been delivered and the fetal membranes have separated. Pain during and after parturition, increased abdominal pressure, retained fetal membranes, and high-oestrogen feed ingredients may contribute to conditions that predispose animals to prolapse. These factors can increase pressure on the reproductive tract. Animals with uterine prolapse require immediate treatment. If treatment is delayed, the exposed organ may become swollen, contaminated, and damaged, and internal haemorrhage may cause death within a few hours. Most animals experiencing uterine prolapse are reported to be hypocalcaemic. Calcium borogluconate should be administered when indicated by veterinary assessment. Animals that recover after appropriate treatment may become pregnant again without further complications.

Retention of Fetal Membranes

Retention of fetal membranes (RFM) is one of the major postpartum complications in dairy goats. It adversely affects animal health, reproductive efficiency, and overall productivity. RFM is generally defined as failure to expel the fetal membranes within approximately 6–8 hours after parturition. Its incidence varies from 2% to 10%.

Hydrometra or Pseudopregnancy (Cloudburst)

Hydrometra is the accumulation of fluid within the uterine lumen. Its exact aetiology is not fully understood, but it is consistently associated with elevated progesterone levels, which are maintained by a persistent corpus luteum (CL). Associated findings may include cessation of cyclic ovarian activity and an increase in abdominal girth. One possible cause of hydrometra is early embryonic loss.

Pseudopregnancy occurs in two forms:

1. Following mating: The early embryo dies after fertilisation, but the corpus luteum remains functional and continues to maintain the uterus in a pregnancy-like state. The abdomen enlarges, but parturition does not occur at the expected time. In some animals, mammary development may occur without the onset of lactation.
2. Without mating: Ovarian cyclic activity ceases, but there is no detectable embryo. At the end of the cycle, the affected animal may expel a large quantity of uterine fluid.

The term cloudburst is used to describe the discharge of a large amount of fluid from the uterus following the resolution of pseudopregnancy. The abdominal enlargement subsequently subsides. In some cases, a few kids may be found, although these may have been missed during the initial examination. After the expected kidding date has passed, ovarian cyclic activity may cease without obvious hydrometra. At the end of the cycle, the affected animal may undergo a bloody discharge. A doe that does not return to oestrus after mating during the breeding season should be examined for possible pseudopregnancy. Treatment with $\text{PGF}_{2\alpha}$ may be indicated under veterinary supervision. Hydrometra can be distinguished from normal pregnancy using B-mode ultrasonography. In hydrometra, the uterus contains fluid but lacks an identifiable embryo or placentomes.

Ovarian Cystic Disease

Ovarian cysts measuring approximately 1.2–3.7 cm have been reported in goats. Cystic ovarian disease has been described in dairy goats, particularly where enlarged follicles and fluid-filled ovarian structures are present.

Abortion

Stress-induced abortion is among the most common causes of pregnancy loss. It may occur due to rough handling, inadequate feeding, transportation, and adverse climatic conditions, particularly during the fourth month of pregnancy. Abortion may result in the birth of either a live or a dead fetus. Habitual abortion may be associated with an inherited predisposition to premature activation of the maternal hypothalamic–pituitary–adrenal axis, which initiates parturition earlier than expected. Such premature births frequently involve dead, weak, or underdeveloped kids.

Nutritional causes

Deficiencies of vitamin A, certain minerals such as manganese and iodine, and dietary energy may reduce reproductive performance. If these deficiencies persist for a prolonged period, they may contribute to abortion.

Infectious Causes of Abortion

Caprine arthritis-encephalitis

Caprine arthritis-encephalitis virus infection may be associated with reproductive problems, although abortion is not considered its characteristic manifestation. The text identifies it among the infectious conditions relevant to goat health.

Listeriosis

Encephalitis caused by *Listeria monocytogenes* infection is relatively common in goats. The organism may also cause late-term abortion or stillbirth. Other infectious diseases that may cause abortion include: Brucellosis, Q fever, Salmonellosis, Toxoplasmosis.

Chlamydial abortion (enzootic abortion)

Chlamydial abortion occurs due to infection with *Chlamydia abortus*. Abortion generally occurs during the final four weeks of pregnancy.

Listeriosis and other bacterial infections

In addition to causing encephalitis, *Listeria monocytogenes* may cause abortion in late pregnancy and the birth of dead kids. Other bacterial infections can also contribute to reproductive losses.

Aspergillosis

Aspergillosis may occur as a systemic fungal disease. It can cause abortion during late pregnancy, sometimes without obvious clinical signs. Affected animals may abort or deliver dead or weak kids, and abnormal postpartum discharge may occur.

References

1. Dhama, K., Karthik, K., Tiwari, R., Shabbir, M. Z., Malik, S. V. S., Singh, R., Chaicumpa, W., & Singh, R. K. (2015). Listeriosis in animals, its public health significance (food-borne zoonosis) and advances in diagnosis and control: A comprehensive review. *Veterinary Quarterly*, 35(4), 211–235.
2. Smith, M. C., & Sherman, D. M. (2009). *Goat medicine* (2nd ed.). Wiley-Blackwell