AGRI MAGAZINE
(International E-Magazine for Agricultural Articles)
Volume: 03, Issue: 11 (November, 2026)
Available online at http://www.agrimagazine.in
©Agri Magazine, ISSN: 3048-8656

Nutrient Mobility: Understanding the Hidden Relationship Between Soil and Plants

*Kotha Shiva Prasad[1], Saketi Radha Krishna[2] and Singireddy PrabhumitraReddy[3]
[1]Ph.D. Scholar, Department of Agricultural Meteorology, IGKV, Raipur, India
[2]Ph.D. Scholar, Department of Vegetable Science, IGKV, Raipur, India
[3]Ph.D. Scholar, Department of Agronomy, IGKV, Raipur, India
*Corresponding Author's email: shiva.prasad0836@gmail.com

Healthy crop growth depends not only on the presence of essential nutrients in the soil but also on their availability to plant roots. A soil may contain considerable quantities of nutrients, yet plants may experience nutrient deficiencies if those nutrients cannot reach the roots in an available form. Similarly, the ability of plants to absorb nutrients depends on their root development, distribution and capacity to explore the surrounding soil. Understanding these interactions is essential for improving soil fertility management and crop productivity. In 1954, Roger H. Bray proposed an important concept known as the nutrient mobility concept of soil-plant relationships. This concept explained how the movement of nutrients in soil and the growth of plant roots together determine nutrient availability and crop response.
The scientific understanding of soil fertility has developed through several important concepts.

Liebig's Law of the Minimum

Justus von Liebig proposed the Law of the Minimum, which states that plant growth is controlled by the nutrient available in the least adequate amount relative to the plant's requirement. For example, a crop may have sufficient nitrogen, potassium and other nutrients, but if phosphorus is deficient, crop growth and yield will be restricted by phosphorus. Even when other nutrients are available in sufficient quantities, they cannot fully compensate for the deficiency of the limiting nutrient. This principle highlights the importance of identifying and correcting the most limiting nutrient for successful crop production.

Mitscherlich's Law of Diminishing Yield Increment

Mitscherlich introduced the concept of diminishing yield increments, explaining that crop yield increases with increasing nutrient application, but the additional yield obtained from each successive application gradually decreases. For example, the first application of nitrogen fertilizer may produce a considerable increase in crop growth and yield. Further applications may continue to increase yield, but the additional benefit becomes progressively smaller. This concept helps explain why excessive fertilizer application does not always result in proportional increases in crop yield.

Mitscherlich-Baule Percentage Sufficiency Concept

The Mitscherlich-Baule concept proposed that the final yield depends on the combined percentage sufficiency of individual nutrients.
For example, if potassium is sufficient for 90% of the attainable yield and phosphorus is sufficient for 80%, the combined percentage sufficiency would be:
$90 \times 80 / 100 = 72\%$
Thus, the crop may produce approximately 72% of its attainable yield under the conditions considered. However, Bray pointed out that the assumptions behind these concepts cannot

explain every soil-plant relationship. He therefore introduced nutrient mobility as an important factor influencing nutrient availability and plant growth.

What is Nutrient Mobility?

Nutrient mobility refers to the ability of nutrients to move through the soil towards plant roots and become available for absorption. When roots absorb nutrients from the surrounding soil, the nutrient concentration near the root surface decreases. Nutrients from nearby soil regions may then move towards the roots through different processes. At the same time, plant roots grow and extend into new soil regions to explore additional sources of nutrients. According to Bray, nutrient movement and root growth are complementary processes that together determine the ability of plants to obtain nutrients. In simple terms, some nutrients move towards the roots, while for other nutrients, roots must grow towards the nutrients.

Different Nutrients Have Different Mobility

All nutrients do not behave in the same way in soil. Their movement depends on their chemical form, solubility and interaction with soil particles.

1. Highly Mobile Nutrients

Nitrate nitrogen is an important example of a highly mobile nutrient. It is readily soluble in soil water and is not strongly adsorbed by most soil particles. Therefore, it can move through the soil along with water. Rainfall or irrigation may transport nitrate downwards through the soil profile. As the soil dries, water movement may also occur towards the surface. Because nitrate is relatively mobile, plant roots can obtain it from a comparatively large volume of soil.

2. Relatively Immobile Nutrients

Nutrients such as exchangeable potassium, calcium and magnesium are held on the surfaces of negatively charged clay and organic matter particles. They are present in exchangeable forms and can be released into the soil solution when roots absorb nutrients and disturb the chemical equilibrium. However, their movement through the soil is relatively limited compared with nitrate. Phosphorus is another important example of a relatively immobile nutrient. It may be strongly adsorbed by soil minerals or present in sparingly soluble forms. Therefore, phosphorus availability is often closely associated with the soil immediately surrounding the roots.

This difference in mobility explains why plants require different mechanisms to obtain different nutrients.

Root Sorption Zones: How Plant Roots Obtain Nutrients

One of the most important contributions of Bray's concept is the explanation of two types of root sorption zones.

1. Root System Sorption Zone

The root system sorption zone represents the larger volume of soil explored by the entire root system. Plants obtain relatively mobile nutrients, such as nitrate nitrogen, from this zone. Since these nutrients can move through the soil, roots can absorb them from a relatively large surrounding soil volume. The effectiveness of this zone depends on root development, root distribution and the movement of nutrients through the soil.

2. Root Surface Sorption Zone

The root surface sorption zone is the thin layer of soil immediately adjacent to the root surface or root hairs. Relatively immobile nutrients, such as exchangeable potassium and adsorbed phosphate, are mainly obtained from this zone. Since these nutrients cannot move over long distances, roots must grow and explore fresh soil regions to obtain an adequate supply.

The important difference is that mobile nutrients are obtained from a larger soil volume, whereas relatively immobile nutrients are obtained mainly from the small soil regions immediately surrounding the roots.

How Root Growth Influences Nutrient Absorption

Root growth plays a major role in determining the amount of nutrients absorbed by plants. As roots extend into new soil regions, they explore additional areas containing nutrients. This is particularly important for relatively immobile nutrients because their movement towards roots is limited. A well-developed root system with greater root length and distribution can explore a larger soil volume and improve access to nutrients. However, simply increasing root growth does not guarantee that every nutrient will be absorbed in sufficient quantities. The chemical form and mobility of the nutrient must also be considered. Thus, nutrient availability depends on both the characteristics of the soil and the ability of the plant to explore it.

Plant Competition and Nutrient Mobility

When plants are grown close together, their root systems may overlap and compete for nutrients.

Bray explained that competition between neighbouring plants can occur in three stages.

Stage 1: No competition

When plants are sufficiently far apart, their root systems do not overlap. Each plant can explore its own soil volume without significant competition for nutrients.

Stage 2: Competition for mobile nutrients

As plants are planted closer together, their root system sorption zones begin to overlap. At this stage, plants may compete for relatively mobile nutrients obtained from the larger soil volume.

Stage 3: Competition for both mobile and immobile nutrients

When plants are grown even closer, their root surface sorption zones may also overlap. Consequently, competition occurs for both relatively mobile and relatively immobile nutrients. This explains why plant spacing and planting density influence nutrient requirements and crop performance.

Bray also emphasized that competition for mobile nutrients may become important earlier than competition for relatively immobile nutrients.

Influence of Plant Species and Rooting Habits

Different plant species and varieties have different root architectures and rooting habits. Some plants develop dense and extensive root systems, while others have relatively shallow or less extensive roots. These differences influence their ability to explore soil and obtain nutrients. Plants with extensive root systems may access a larger volume of soil and obtain more mobile nutrients. However, their ability to obtain relatively immobile nutrients also depends on how effectively their roots and root hairs explore fresh soil regions. Therefore, nutrient requirements may differ among plant species and varieties because of differences in their root systems.

Influence of Growth Stage on Nutrient Mobility

Nutrient requirements and root absorption capacity change during different stages of plant growth. At certain growth stages, the root system may not be sufficiently developed to absorb nutrients according to the plant's requirements. This can influence the availability of nutrients, particularly those that are relatively immobile. Bray suggested that the growth stage at which a plant develops its greatest nutrient requirement is important in determining the soil concentration of relatively immobile nutrients needed for maximum yield. In general, an adequate supply of nutrients during early growth is particularly important because deficiencies of relatively immobile nutrients at early stages may restrict subsequent plant development.

Nutrient Distribution and Soil Fertility Patterns

The distribution of nutrients within soil also influences their availability to plants. An important observation made by Bray was that a uniformly distributed quantity of a relatively immobile nutrient may not always provide the highest availability to roots. When nutrients

are concentrated in particular soil regions, root systems may obtain them more effectively than when the same quantity is distributed uniformly at a lower concentration. This means that nutrient availability depends not only on the total quantity present in the soil but also on its distribution and concentration around the roots. For example, the localized placement of relatively immobile nutrients such as phosphorus can improve their accessibility to developing roots under suitable soil conditions.

Influence of Soil and Seasonal Conditions

Soil properties and seasonal conditions influence nutrient movement and plant growth. Factors such as soil texture, clay content, organic matter, rainfall and soil moisture affect nutrient availability and mobility. For example, nitrate may move readily through soil water, whereas potassium and phosphorus are more strongly influenced by soil adsorption and exchange reactions. Similarly, rainfall and seasonal conditions influence plant size and root development. A favourable growing season may promote larger plants with more extensive root systems, allowing them to explore a greater soil volume. These factors demonstrate that nutrient requirements are not determined by nutrient concentration alone. The physical and chemical properties of soil and the growing environment must also be considered.

Integrating Liebig's and Mitscherlich's Concepts Through Nutrient Mobility

Bray's nutrient mobility concept provides a framework for understanding the conditions under which Liebig's and Mitscherlich's concepts operate. Relatively mobile nutrients, such as nitrate nitrogen, can be absorbed from a large volume of soil explored by the root system. Their availability and requirement are strongly associated with the crop's yield potential and nutrient composition. On the other hand, relatively immobile nutrients, such as phosphorus and exchangeable potassium, are obtained mainly from the soil immediately surrounding the roots. Their availability depends greatly on root growth, root distribution and nutrient concentration in the root surface sorption zone. Thus, the nutrient mobility concept explains why the limiting nutrient principle and percentage sufficiency concept may apply under particular conditions rather than universally.

Nutrient Mobility and Scientific Fertility Management

Understanding nutrient mobility has important implications for soil fertility management.

- **Fertilizer placement:** Relatively immobile nutrients should be placed where developing roots can access them effectively.
- **Root development:** Healthy and extensive root systems improve the exploration of soil and nutrient acquisition.
- **Plant spacing:** Appropriate spacing can reduce excessive competition between neighbouring plants.
- **Soil testing:** Nutrient concentration alone should be interpreted along with soil properties and nutrient availability.
- **Crop-specific management:** Fertilizer recommendations should consider crop species, rooting habits and growth stages.
- **Water management:** Proper irrigation and drainage help regulate nutrient movement and reduce nutrient losses.

These practices help improve nutrient-use efficiency and support better crop growth.

Conclusion

The nutrient mobility concept proposed by Roger H. Bray provides a valuable understanding of the relationship between soil nutrients and plant roots. It emphasizes that the availability of nutrients depends not only on their quantity in soil but also on their chemical form, mobility, distribution and the ability of roots to explore the soil. Highly mobile nutrients are generally obtained from a larger volume of soil, whereas relatively immobile nutrients are absorbed mainly from the small soil regions surrounding the roots. Root growth, plant spacing, growth stage, soil properties and seasonal conditions further influence these processes.

Therefore, successful soil fertility management requires an understanding of both nutrient behaviour in soil and plant root development. The concept reminds us that **soil fertility is not simply about how much nutrient is present, but also about how effectively plants can reach and absorb it.**

References

1. Bray RH. (1938). New concepts on the chemistry of soil fertility. *Soil Science Society of America Proceedings*, 2, 175-179.
2. Bray RH. (1942). Ionic competition in base-exchange reactions. *Journal of the American Chemical Society*, 64, 954-963.
3. Bray RH. (1944). Soil-plant relations: I. Soil Sci., 58, 305-324.
4. Bray RH. (1944). Soil-plant relations: II. *Soil Science*, 60, 463-473.
5. Deturk EE. (1942). The phosphate fertilizers problem. *Illinois Agricultural Experiment Station Bulletin*, 484, 561-562.
6. Jenny H and Overstreet R. (1939). Cation interchange between plant roots and soil colloids. *Soil Science*, 47, 257-272.
7. Kurtz LT, DeTurk EE and Bray RH. (1946). Phosphate adsorption by Illinois soils. *Soil Science*, 61, 111-124.
8. Liebig, J. von. (1855). *Die Grundsätze der Agriculturchemie*. Braunschweig.
9. Mitscherlich, E. A. (1909). Das Gesetz des Minimums und das Gesetz des abnehmenden Bodenertrages. *Landwirtschaftliches Jahrbuch*, 38, 537–552.
10. Spillman WJ and Lang E. (1924). *The Law of Diminishing Returns*. World Book Company.